

# Quadrature in meshless methods for general elliptic problems with natural boundary conditions

Qinghui Zhang <sup>\*</sup>      Uday Banerjee <sup>†</sup>

## Abstract

In this paper, we explore the effect of numerical integration on the meshless method used to approximate the solution of an elliptic partial differential equation with non-constant coefficients with Neumann boundary conditions. We considered meshless methods with shape functions that reproduce polynomials of degree  $k \geq 1$ . We have obtained an estimate for the energy norm of the error in the approximate solution from the meshless method under the presence of numerical integration. This result was established under the assumption that the numerical integration rule satisfied a certain discrete Green's formula, which is not problem dependent, i.e., does not depend on the non-constant coefficients of the problem. We have also derived numerical integration rules in two dimensions satisfying the discrete Green's formula.

**Keywords:** PDE with non-constant coefficients, Galerkin methods, meshless methods, quadrature, numerical integration, error estimates

## 1 Introduction

Since last 20 years, a lot of progress has been made in the development of the Meshless Method (MM), and it has been applied to solve complicated engineering problems (see e.g., [1, 2, 3, 8, 19, 22, 25]). The implementation of MM does not require a mesh, but unlike standard the Finite Element Method (FEM), the shape functions used in the MM are not piecewise polynomials. This feature of the shape functions poses a serious challenge in the use of numerical integration

---

<sup>\*</sup>Department of Scientific Computing and Computer Applications, Sun Yat-Sen University, Guangzhou, 510275, P. R. China. E-mail address: zhqingh@mail2.sysu.edu.cn. This research was partially supported by the NSF Grant # DMS-0610778, the China Scholarship Council through the State Scholarship Program, and the National Science Foundation of China grant 10771224

<sup>†</sup>Department of Mathematics, 215 Carnegie, Syracuse University, Syracuse, NY 13244. E-mail address: banerjee@syr.edu. <http://banerjee.syr.edu>. This research was partially supported by the NSF Grant # DMS-0610778.

to compute the elements of the stiffness matrix, the mass matrix, and the load vector.

From the very beginning of the development of the MM, it has been recognized that numerical integration is a serious issue in MM and has been addressed in various engineering papers [9, 18, 20, 7, 13, 14, 5, 16, 17, 11, 12]. Several approaches to implement numerical integration have been proposed in the literature; we refer to Section 3 of [6] for a brief review of these approaches. A mathematical analysis of the effect of numerical integration was first reported in [5], where it was shown that the approximate solution obtained from the MM, using standard numerical integration, may not converge. It was also shown that if the stiffness matrix (numerically computed with quadrature) satisfies a row sum condition then the error in the approximate solution (in energy norm) is  $O(h + \eta)$ , where  $h$  is the discretization parameter and the parameter  $\eta$  indicates the accuracy of the underlying numerical integration. Thus with  $\eta = O(h)$ , the MM with numerical integration yields the optimal order of convergence. However, the analysis presented in [5] was restricted to the shape functions of the MM that reproduced polynomials of degree  $k = 1$  and could not be extended for  $k > 1$ .

Another analysis of the effect of numerical integration on MM was presented later in [6], where the quadrature is required to satisfy a discrete Green's formula. This analysis is valid for the MMs, where the shape functions reproduced polynomials of degree  $k \geq 1$ . It was shown that the energy-norm of the error in the approximate solution obtained from MM is  $O(h^{k-1}(h + \eta))$ , and optimal order of convergence is obtained with  $\eta = O(h)$ . However in [6], the MM was used to approximate the solution of a Neumann problem with constant coefficient with no lower order term. We further note that a direct application of the ideas in [6] to the situation, where MM is applied to a problem with non-constant coefficient, required the quadrature to be problem dependent, e.g., dependent on the non-constant coefficients of the problem.

In this paper we extend the analysis in [6] to study the effect of numerical integration, when the MM is used to approximate the solution of a Neumann problem with non-constant coefficients including the lower order term. We require the quadrature to satisfy an extended version of the discrete Green's formula, which is not problem dependent. We show that the energy-norm of the error in the approximate solution obtained from MM with quadrature is  $O(h^{k-1}(h + \eta))$ . For a Neumann problem with no lower order term, we mention that the condition on the quadrature required in this paper is same as the condition required in [6] for  $k = 1$ . However for  $k > 1$ , the situation is different; a quadrature satisfying the condition proposed in this paper automatically satisfies the condition required in [6], but not vice versa. In this paper, we have also investigated the possibility of using different numerical integration rules to compute the elements of the stiffness matrix, the mass matrix, and the load vector, which was not done in [6]. Moreover, we have derived numerical integration rules, satisfying the extended discrete Green's formula, in two dimensions for  $k = 1$  and  $k = 2$  in this paper; numerical integration rules only for  $k = 1$  in one dimension was presented in [6].

The outline of the paper is as follows: In Section 2, we presented the notations and the elliptic Neumann model problem with non-constant coefficients. The MM and the various properties of the associated finite dimensional space is given in Section 3. In Section 4, we defined the MM with numerical integration and listed the assumptions imposed on the numerical integration rule. The effect of numerical integration on the energy norm of the error in the approximate solution, obtained from the MM, has been investigated in Section 5. Our main analytical result, Theorem 5.5, have also been presented in this section. In Section 6, we derived numerical integration rules, in 2-dimensions, that satisfy the main assumption given in Section 4. We presented some numerical examples in Section 7 that shows the effect of the numerical integration on the energy norm of the error in the approximate solution.

## 2 Preliminaries and model problem

Let  $\mathbb{N}$  the set of all positive integers. For a domain  $D \subset \mathbb{R}^d$ , an integer  $m \in \mathbb{N} \cup \{0\}$ , and  $p \in \mathbb{N} \cup \infty$ , we denote the usual Sobolev space by  $W^{m,p}(D)$  with the norm  $\|\cdot\|_{W^{m,p}(D)}$  and semi-norm  $|\cdot|_{W^{m,p}(D)}$ . We will only consider  $p = 2, \infty$  in this paper. The Sobolev space  $W^{m,p}(D)$  will be represented by  $H^m(D)$  in the case  $p = 2$  and by  $L_p(D)$  in the case  $m = 0$ . Likewise, for a hypersurface  $\partial D$  in  $\mathbb{R}^d$ , we will use the space  $L_p(\partial D)$  equipped with the norm  $\|\cdot\|_{L_p(\partial D)}$ .

Let  $V$  be a normed linear space. We define  $\tilde{V}$  to be the product space  $V^d$ , where  $\tilde{v} = [v_i]_{i=1}^d \in \tilde{V}$  is a vector-valued function with its component  $v_i \in V$ ,  $i = 1, 2, \dots, d$ . When  $V = W^{m,p}(D)$  or  $L_p(\partial D)$ , the associated norm of  $\tilde{V}$  is defined by  $\|\tilde{v}\|_V = (\sum_{i=1}^d \|v_i\|_V^p)^{\frac{1}{p}}$  in the case  $1 \leq p < \infty$  and  $\|\tilde{v}\|_V = \max\{\|v_i\|_V : i = 1, 2, \dots, d\}$  in the case  $p = \infty$ ; the semi-norm  $|\tilde{v}|_V$  (for  $V = W^{m,p}(D)$ ) is defined by using  $|v_i|_V$  instead of  $\|v_i\|_V$  in the above definition.

A domain  $D$  is star-shaped with respect to a ball  $B \subset D$  if, for all  $x \in D$ , the closed convex hull of  $\{x\} \cap B$  is a subset of  $D$ . Let  $\rho_{\max} = \sup\{\rho : D \text{ is star-shaped with respect to a ball of radius } \rho\}$ , then the *chunkiness parameter* of  $D$  is defined by

$$\gamma_D = \frac{\text{diam}(D)}{\rho_{\max}}.$$

Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain with Lipschitz continuous boundary  $\Gamma = \partial\Omega$ . For the model problem, we consider the Neumann problem

$$\begin{aligned} \mathcal{L}u \equiv -\nabla \cdot (A \nabla u) + cu &= f, & \text{in } \Omega \\ A \nabla u \cdot \vec{n} &= g, & \text{on } \Gamma \end{aligned} \quad (2.1)$$

where  $A(x) = \{a_{ij}(x)\}_{1 \leq i, j \leq d}$  is a symmetric matrix,  $a_{ij} \in C^k(\bar{\Omega})$ ,  $c \in C(\bar{\Omega})$ ,  $f \in L_2(\Omega)$ ,  $g \in L_2(\Gamma)$  and  $\vec{n}$  is the outward unit normal vector to  $\Gamma$ . We assume that there is a constant  $\alpha > 0$  such that

$$\sum_{i,j=1}^d u_i a_{ij}(x) u_j \geq \alpha \sum_{i=1}^d u_i^2, \forall u \in \mathbb{R}^d \text{ and } c(x) \geq \alpha, \quad \forall x \in \Omega. \quad (2.2)$$

We note that for  $v \in H^2(\Omega)$ ,  $a\nabla v$  is a vector-valued function, but for simplicity of notation, we do not put a tilde over it.

The associated variational formulation of (2.1) is given by

$$\begin{aligned} &\text{Find } u \in H^1(\Omega) \text{ such that} \\ &B(u, v) = L(v), \quad \forall v \in H^1(\Omega) \end{aligned} \quad (2.3)$$

where

$$B(u, v) \equiv B_1(u, v) + B_0(u, v), \quad L(v) \equiv \int_{\Omega} f v \, dx + \int_{\Gamma} g v \, ds$$

and

$$B_1(u, v) \equiv \int_{\Omega} A \nabla u \cdot \nabla v \, dx, \quad B_0(u, v) \equiv \int_{\Omega} c u v \, dx$$

It is clear that the bilinear form  $B(\cdot, \cdot)$  is continuous and coercive on  $H^1(\Omega) \times H^1(\Omega)$ , and it is well known [10] that the variational problem (2.3) has a unique solution.

### 3 Meshless methods

The meshless method to approximate the solution of the variational problem (2.3) is a Galerkin method, where the construction of the underlying finite dimensional subspace either does not depend on a mesh, or uses a mesh only minimally. To this end, we consider a one-parameter family of finite dimensional spaces  $V_h \subset H^1(\Omega)$ , given by

$$V_h = \text{span}\{\phi_i^h \in C(\Omega) : i \in N_h\}; \quad N_h \text{ is an index set.}$$

The functions  $\{\phi_i^h\}_{i \in N_h}$  are referred to as *shape functions* and their construction either does not depend on a mesh or depends only minimally. Each  $\phi_i^h$  has compact support and we let  $\omega_i^h$  denote the interior of  $\text{supp } \phi_i^h$  with  $h_i = \text{diam } \omega_i^h$ . We assume that each  $\omega_i^h$  is star-shaped with respect to a ball  $o_i^h \subset \omega_i^h$  and their chunkiness parameters satisfy  $\gamma_{\omega_i^h} \leq C, \forall i \in N_h$ .

Often the shape function  $\{\phi_i^h\}_{i \in N_h}$  are constructed relative to a set of *particles*  $X_h = \{x_i^h : i \in N_h\} \subset \mathbb{R}^d$  and each  $\phi_i^h$  is associated with a particle  $x_i^h$ . When  $\bar{\omega}_i^h \cap \Gamma = \emptyset$ , then the associated particle  $x_i^h \in \omega_i^h$ . But when  $\bar{\omega}_i^h \cap \Gamma \neq \emptyset$ , then the associated particle  $x_i^h$  could be outside  $\Omega$ . We divide the index set  $N_h$  into two disjoint parts,  $N'_h$  and  $N''_h$ , where,

$$N'_h = \{i \in N_h : \partial\omega_i \cap \Gamma \neq \emptyset\} \text{ and } N''_h = \{i \in N_h : \bar{\omega}_i \subset \Omega\}.$$

Now, we make several assumptions on the space  $V_h$ .

**A1:** For  $i \in N_h$ , let  $S_i \equiv \{j \in N_h : \omega_i^h \cap \omega_j^h \neq \emptyset\}$ . We assume that there is a constant  $\kappa$ , independent of  $i$ ,  $j$ , and  $h$ , such that

$$\text{card } S_i \leq \kappa.$$

**Remark 3.1** This property is referred to as the *finite overlap property*. If we let  $S_x = \{j \in N_h : x \in \omega_j^h\}$ , then the finite overlap property implies

$$\text{card } S_x \leq \kappa, \quad \forall x \in \Omega. \quad (3.1)$$

**A2:** There exist positive constants  $C_1$  and  $C_2$ , independent of  $h$  and  $i$ , such that

$$C_1 \leq \frac{h_i}{h} \leq C_2, \quad C_1 h^d \leq |\omega_i| \leq C_2 h^d, \quad \text{and} \quad C_1 h^{d-1} \leq |\bar{\omega}_i \cap \Gamma| \leq C_2 h^{d-1}, \quad (3.2)$$

where  $|\omega_i|$  is the “area” of  $\omega_i$  in  $\mathbb{R}^d$  and  $|\bar{\omega}_i \cap \Gamma|$  is the “length” of  $\bar{\omega}_i \cap \Gamma$  in  $\mathbb{R}^{d-1}$ .

**A3:** The shape functions *reproduce polynomials of degree  $k$* , i.e.,

$$\sum_{i \in N_h} p(x_i^h) \phi_i^h(x) = p(x), \quad \forall p \in \mathcal{P}^k \text{ and } x \in \Omega. \quad (3.3)$$

**A4:** There exists a positive constant  $C$ , independent of  $i$  and  $h$ , such that

$$\|D^\alpha \phi_i^h\|_{L^\infty(\mathbb{R}^d)} \leq C h_i^{-|\alpha|} \text{ for } |\alpha| \leq q \text{ for some } q \geq 1. \quad (3.4)$$

In this paper, we will assume  $q = k + 1$ .

**A5:** There exist positive constants  $C_1, C_2$ , independent of  $h$  and  $i$ , such that for any  $i \in N_h$ ,

$$C_1 \|v\|_{L_2(\omega_i)}^2 \leq h_i^d \sum_{j \in S_i} v_j^2 \leq C_2 \|v\|_{L_2(\omega_i)}^2, \quad (3.5)$$

$$C_1 \|v\|_{L_2(\partial\omega_i \cap \Gamma)}^2 \leq h_i^{d-1} \sum_{j \in S'_i} v_j^2 \leq C_2 \|v\|_{L_2(\partial\omega_i \cap \Gamma)}^2, \quad (3.6)$$

$$C_1 |v|_{H_1(\omega_i)}^2 \leq h_i^{d-2} \sum_{j \in S_i} (v_j - v_i)^2 \leq C_2 |v|_{H_1(\omega_i)}^2 \quad (3.7)$$

where  $v \in V_h$  is of the form  $v = \sum_{i \in N_h} v_i \phi_i^h$  and  $S'_i \equiv \{j \in N'_h : \partial\omega_j \cap (\partial\omega_i \cap \Gamma) \neq \emptyset\} \subset N_h$ .

**Remark 3.2** The finite dimensional space  $V_h$  could be viewed as a generalization of the standard piecewise linear finite element space based on quasi-uniform mesh. In the finite element setting, the shape functions  $\phi_i^h$  are hat functions, the particles  $x_i^h$  are the finite element nodes, and the supports  $\bar{\omega}_i^h$  are the finite element “stars”. The quasi-uniform finite elements satisfy the assumptions **A1** and **A2**, whereas the hat functions satisfy **A3** and **A4** with  $k = 1$ . The inequalities (3.5), (3.6), and (3.7) are also true for piecewise linear finite elements. But finite elements are piecewise polynomials and their construction requires a mesh. ■

**Remark 3.3** Many approaches to construct shape functions for meshless methods are available primarily in the engineering literature; we refer to [8, 9, 16, 21, 24, 23] for details. In all these approaches, the shape functions are not piecewise polynomials and are not available in terms of explicit mathematical formulas that could be easily evaluated. This is the price one pays for avoiding a mesh. For example, in the reproducing kernel particle (RKP) technique, one starts with a weight function  $w(x)$  with compact support such that the origin is in the interior of the support. The shape function  $\phi_i^h$  is sought in the form

$$\phi_i^h(x) = w_i^h(x) \sum_{|\alpha| \leq k} (x - x_i^h)^\alpha b_\alpha^h(x), \quad (\alpha \text{ is a multi-index})$$

where  $w_i^h(x) = w(\frac{x-x_i}{h_i})$ . For each  $x \in \Omega$ ,  $b_\alpha^h(x)$  are chosen so that (3.3) is satisfied, which requires solving a linear system. For details, we refer the reader to [21, 25]. We also note that the shape functions  $\phi_i^h(x)$ , constructed using these approaches, do not satisfy the *Kronecker delta property*, i.e.,  $\phi_i^h(x_j^h) \neq \delta_{ij}$ . We further note that if the weight function  $w \in C^q(\Omega)$ , then the shape functions  $\phi_i^h \in C^q(\Omega)$ . Thus it is easy to construct smooth shape functions, in particular with  $q = k + 1$ , as assumed in assumption **A4**. ■

**Remark 3.4** The inequality (3.5) in assumption **A5** implies the local linear independence of the shape functions  $\{\phi_j : j \in S_i\}$ . The constants  $C_1$  and  $C_2$  appearing in **A5** may depend on the geometry of  $\omega_i$  and  $\omega_j$  with  $j \in S_i$ , but are independent of  $i$  and  $h$ . We note that the assumption **A5** is weaker than a similar assumption used in [6]. ■

**Remark 3.5** Summing the terms of the inequality (3.5) over  $i \in N_h$  and using the assumption **A1** and **A2**, we can obtain

$$C_1 \|v\|_{L_2(\Omega)}^2 \leq h^d \sum_{i \in N_h} v_i^2 \leq C_2 \|v\|_{L_2(\Omega)}^2, \quad \forall v = \sum_{i \in N_h} v_i \phi_i \in V_h \quad (3.8)$$

Similarly, we obtain

$$C_1 \|v\|_{L_2(\Gamma)}^2 \leq h^{d-1} \sum_{i \in N'_h} v_i^2 \leq C_2 \|v\|_{L_2(\Gamma)}^2, \quad \forall v = \sum_{i \in N'_h} v_i \phi_i \in V_h. \quad (3.9)$$

In particular, substituting  $v = 1$  in these two inequalities, we get

$$\begin{aligned} C_1 h^{-d} \leq |N_h| \leq C_2 h^{-d}, \quad C_2 h^{-d} \leq |N''_h| \leq C_2 h^{-d} \\ C_1 h^{-(d-1)} \leq |N'_h| \leq C_2 h^{-(d-1)} \end{aligned} \quad (3.10)$$

These estimates will be used later in the paper. ■

In the rest of the paper, we will suppress the parameter  $h$  for notational clarity and write  $\phi_i$ ,  $x_i$ ,  $\omega_i$ , and  $o_i$  for  $\phi_i^h$ ,  $x_i^h$ ,  $\omega_i^h$ , and  $o_i^h$ , respectively, with the understanding that they depend on  $h$ .

Based on the finite dimensional space  $V_h \subset H^1(\Omega)$ , as described above, *the meshless method* to approximate the solution of (2.3) is given by

$$\begin{aligned} &\text{Find } u_h \in V_h \text{ such that} \\ &B(u_h, v_h) = L(v_h), \quad \forall v_h \in V_h \end{aligned} \quad (3.11)$$

The approximation of the exact solution  $u \in H^1(\Omega)$  by the solution  $u_h \in V_h$  of (3.11) depends on the approximation property of the space  $V_h$ , which has been studied in [21, 25]. But in these studies, the set of particles,  $X_h$ , has been assumed to be in  $\Omega$ , which may give rise to boundary layer in the error as indicated in ([4]). This is precisely the reason some of the particles have been allowed to be outside  $\Omega$  in this paper, as well as in [3, 5, 6]. But the approximation result for  $V_h$  remains the same as in [21, 25], even when some of the particles are allowed to outside  $\Omega$ ; only the analysis requires slight modification based on an extension result. For completeness, we present the modified analysis in this paper.

For a function  $u \in W^{k+1, \infty}(\mathbb{R}^d) \cap C(\mathbb{R}^d)$ , we define its  $V_h$ -“interpolant” on  $\Omega$  by  $\mathcal{I}_h u$ , given by

$$\mathcal{I}_h u(x) = \sum_{i \in N_h} u(x_i) \phi_i(x), \quad x \in \Omega. \quad (3.12)$$

It is clear from the reproducing property (3.3) that  $\mathcal{I}_h p(x) = p(x)$ , for  $x \in \Omega$  and  $p \in \mathcal{P}^k$ . Strictly speaking,  $\mathcal{I}_h$  is not an interpolation operator since  $\mathcal{I}_h u(x_j) \neq u(x_j)$  for  $x_j \in X_h$ .

When  $u$  is defined only on  $\Omega$ , the interpolant  $\mathcal{I}_h u$  is undefined as some of the particles  $x_i$  may be outside  $\Omega$ . To address this issue, we use the well-known extension theorem (see [10, 26]), which provides us with an extension operator

$$\begin{aligned} E : L_2(\Omega) &\rightarrow L_2(\mathbb{R}^d), \\ u &\mapsto \bar{u} \equiv Eu \end{aligned}$$

such that

$$\bar{u}(x) = u(x), \text{ for } x \in \Omega \text{ and } \|Eu\|_{W^{k+1, \infty}(\mathbb{R}^d)} \leq C \|u\|_{W^{k+1, \infty}(\Omega)} \quad (3.13)$$

where constant  $C$  is independent of  $u \in L_2(\Omega)$ . We now define the  $V_h$ -interpolant of  $u \in W^{k+1, \infty}(\Omega) \cap C(\Omega)$  by  $\mathcal{I}_h u(x) \equiv \mathcal{I}_h \bar{u}(x)$ , for  $x \in \Omega$ . We now present an interpolation result for  $V_h$ , which indicates the approximation property of  $V_h$ .

**Theorem 3.1** *Let  $u \in W^{k+1, \infty}(\Omega) \cap C(\Omega)$  and  $\mathcal{I}_h u$  be the  $V_h$ -interpolant of  $u$ . Then there is a positive constant  $C$ , independent of  $h$ , such that*

$$\|u - \mathcal{I}_h u\|_{W^{l, p}(\Omega)} \leq Ch^{k+1-l} \|u\|_{W^{k+1, \infty}(\Omega)} \quad \forall 0 \leq l \leq k+1 \text{ and } p \geq 1. \quad (3.14)$$

*Proof.* For  $i \in N_h$ , let  $\hat{\omega}_i$  be the smallest ball containing the set  $\cup_{j \in S_i} \omega_j$ . Consider  $Q_i^{k+1} E u(x)$ , the Taylor polynomial of degree  $k$  (i.e., of order  $k+1$ ) of

$Eu$  averaged over the ball  $\hat{\omega}_i$  (see the definition 4.1.3 in [10]). Then from the Lemma 4.3.8 of [10] and assumption **A1**, we have

$$\|Eu - Q_i^{k+1}Eu\|_{W^{l,\infty}(\hat{\omega}_i)} \leq Ch^{k+1-l}|Eu|_{W^{k+1,\infty}(\hat{\omega}_i)} \quad \forall 0 \leq l \leq k+1. \quad (3.15)$$

The constant  $C$  depends on  $\kappa$  and the chunkiness parameter of  $\hat{\omega}_i$ , which is 1, and thus  $C$  is independent of  $i$ .

For  $x \in \omega_i$ , we note that  $\phi_j(x) = 0$  for  $j \notin S_i$ . Therefore, for  $x \in \omega_i$ , we have

$$\begin{aligned} u(x) - \mathcal{I}_h u(x) &= Eu(x) - Q_i^{k+1}Eu(x) + Q_i^{k+1}Eu(x) \\ &\quad - \sum_{j \in S_i} [Eu(x_j) - Q_i^{k+1}Eu(x_j)]\phi_j(x) \\ &\quad - \sum_{j \in S_i} [Q_i^{k+1}Eu(x_j)]\phi_j(x) \\ &= Eu(x) - Q_i^{k+1}Eu(x) - \sum_{j \in S_i} [Eu(x_j) - Q_i^{k+1}Eu(x_j)]\phi_j(x), \end{aligned}$$

where we used (3.3) with  $p(x) = Q_i^{k+1}Eu(x)$ . Therefore, from (3.15) and assumptions **A4**, **A1**, **A2** we get

$$\begin{aligned} \|u - \mathcal{I}_h u\|_{W^{l,\infty}(\omega_i)} &\leq \|Eu - Q_i^{k+1}Eu\|_{W^{l,\infty}(\hat{\omega}_i)} \\ &\quad + \sum_{j \in S_i} \|Eu - Q_i^{k+1}Eu\|_{W^{0,\infty}(\hat{\omega}_i)} \|\phi_j\|_{W^{l,\infty}(\hat{\omega}_i)} \\ &\leq Ch^{k+1-l} \|Eu\|_{W^{k+1,\infty}(\hat{\omega}_i)} + \sum_{j \in S_i} Ch^{k+1} \|Eu\|_{W^{k+1,\infty}(\hat{\omega}_i)} h^{-l} \\ &\leq Ch^{k+1-l} \|Eu\|_{W^{k+1,\infty}(\hat{\omega}_i)}, \end{aligned} \quad (3.16)$$

where  $C$  may depend on  $\kappa$ . Thus we immediately get

$$\|u - \mathcal{I}_h u\|_{W^{l,p}(\omega_i)} \leq Ch^{\frac{d}{p}} h^{k+1-l} \|Eu\|_{W^{k+1,\infty}(\hat{\omega}_i)}. \quad (3.17)$$

Finally, using (3.16), (3.17), the assumption **A1**, and (3.10), we get

$$\begin{aligned} \|u - \mathcal{I}_h u\|_{W^{l,\infty}(\Omega)} &\leq \sup_{i \in N_h} \|u - \mathcal{I}_h u\|_{W^{l,\infty}(\omega_i)} \leq Ch^{k+1-l} \sup_{i \in N_h} \|Eu\|_{W^{k+1,\infty}(\hat{\omega}_i)} \\ &\leq Ch^{k+1-l} \|Eu\|_{W^{k+1,\infty}(\mathbb{R}^d)} \leq Ch^{k+1-l} \|u\|_{W^{k+1,\infty}(\Omega)} \end{aligned}$$

and

$$\begin{aligned} \|u - \mathcal{I}_h u\|_{W^{l,p}(\Omega)} &\leq \left[ \sum_{i \in N_h} \|u - \mathcal{I}_h u\|_{W^{l,p}(\omega_i)}^p \right]^{\frac{1}{p}} \\ &\leq Ch^{\frac{d}{p}} h^{k+1-l} \|\bar{u}\|_{W^{k+1,\infty}(\mathbb{R}^d)} |N_h|^{\frac{1}{p}} \\ &\leq Ch^{k+1-l} \|u\|_{W^{k+1,\infty}(\Omega)}, \end{aligned}$$

which gives the desired result.  $\square$



**Remark 3.6** We note that Theorem 3.1 holds for  $u \in W^{k+1,p}(\Omega)$ ,  $1 < p < \infty$ , provided  $k+1 > d/p$  ( $k+1 \geq d$  when  $p = 1$ ). Also for a given  $l$ , we only need  $q = l$  in assumption A4 (instead of  $q = k+1$ ). ■

Now, with Lax-Milgram Theorem, Céa' Theorem [10] and (3.14), the following approximation result for the meshless method with exact integration is immediate:

**Theorem 3.2** *There is a unique solution  $u_h \in V_h$  of the variational problem (3.11) satisfying*

$$\|u - u_h\|_{H^1(\Omega)} \leq Ch^k \|u\|_{W^{k+1,\infty}(\Omega)}, \quad (3.18)$$

where  $C$  is independent of  $h$ .

Another consequence of the interpolation error estimate (3.14) in Theorem 3.1 is

$$\begin{aligned} \|\mathcal{I}_h u\|_{W^{k+1,\infty}(\Omega)} &\leq \|u\|_{W^{k+1,\infty}(\Omega)} + \|u - \mathcal{I}_h u\|_{W^{k+1,\infty}(\Omega)} \\ &\leq C \|u\|_{W^{k+1,\infty}(\Omega)}, \end{aligned} \quad (3.19)$$

which will be used later in this paper. This is the reason we required  $q = k+1$  in assumption **A4**.

## 4 The meshless method with numerical integration

In this section, we will present the meshless method with numerical integration (also referred to as quadrature). We will also state the assumptions imposed on the underlying numerical integration rule and discuss them.

To motivate the quadrature in MM, we write the solution  $u_h$  of (3.11) as  $u_h = \sum_{j \in N_h} c_j \phi_j$ . Then the coefficients  $\{c_j\}_{j \in N_h}$  can be determined uniquely from the linear system

$$\sum_{j \in N_h} (\gamma_{ij} + \sigma_{ij}) c_j = l_i, \quad \forall i \in N_h,$$

where

$$\gamma_{ij} \equiv B_1(\phi_j, \phi_i) = \int_{\Omega} A \nabla \phi_j \cdot \nabla \phi_i \, dx = \int_{\omega_i} A \nabla \phi_j \cdot \nabla \phi_i \, dx,$$

$$\sigma_{ij} \equiv B_0(\phi_j, \phi_i) = \int_{\Omega} c \phi_j \phi_i \, dx = \int_{\omega_i} c \phi_j \phi_i \, dx,$$

and

$$l_i \equiv L(\phi_i) = \int_{\Omega} f \phi_i \, dx + \int_{\Gamma} g \phi_i \, ds = \int_{\omega_i} f \phi_i \, dx + \int_{\partial \omega_i \cap \Gamma} g \phi_i \, ds;$$

we recall that the shape function  $\phi_i$  has compact support  $\bar{\omega}_i$ . We mention that  $\omega_i \cap \omega_j$  can also be used as the domain of integration in the definition of  $\gamma_{ij}$  and  $\sigma_{ij}$ , since the shape function  $\phi_j$  has compact  $\bar{\omega}_j$ . Consequently, the matrices  $\{\gamma_{ij}\}$  and  $\{\sigma_{ij}\}$  are symmetric. We have used  $\omega_i$  (instead of  $\omega_i \cap \omega_j$ ) in the definition of  $\gamma_{ij}$  and  $\sigma_{ij}$  to motivate the numerical integration scheme in this paper. The integrals  $\gamma_{ij}$ ,  $\sigma_{ij}$ ,  $\int_{\omega_i} f \phi_i dx$  and  $\int_{\partial\omega_i \cap \Gamma} g \phi_i ds$  have to be computed numerically using numerical integration formulas on  $\omega_i$ ,  $i \in N_h$  and on  $\partial\omega_i \cap \Gamma$ ,  $i \in N'_h$ . Let

$$\gamma_{ij}^* \equiv B_1^*(\phi_j, \phi_i) \equiv \int_{\omega_i}^s A \nabla \phi_j \cdot \nabla \phi_i dx, \quad \sigma_{ij}^* \equiv B_0^*(\phi_j, \phi_i) \equiv \int_{\omega_i}^m c \phi_j \phi_i dx,$$

and

$$l_i^* \equiv \int_{\omega_i}^l f \phi_i dx + \int_{\partial\omega_i \cap \Gamma} g \phi_i ds,$$

where  $f_{\omega_i}^s$ ,  $f_{\omega_i}^m$ , and  $f_{\omega_i}^l$  denote the numerical integration rules, defined on  $\omega_i$ , to approximate the entries of the stiffness matrix, mass matrix, and the load vector (only the volume integrals), respectively;  $f_{\partial\omega_i \cap \Gamma}$  is the numerical integration rule to approximate the “boundary integral” in the elements of the load vector.

We note that for a given  $i \in N_h$ , we use the same quadrature rule  $f_{\omega_i}^s$  to compute  $\gamma_{ij}^*$  for each  $j \in S_i$  (recall the definition of  $S_i$  in assumption **A1** in Section 3); similarly, the same quadrature rule  $f_{\omega_i}^m$  is used to compute  $\sigma_{ij}^*$  for  $j \in S_i$ . But the quadrature rules  $f_{\omega_i}^s$  and  $f_{\omega_i}^m$  could possibly be different, i.e., different quadrature rules could be used to approximate the integrals in the stiffness matrix and the mass matrix. The idea of using possibly different quadrature rules to compute the stiffness and mass matrix was not considered in [6].

**Remark 4.1** It is easy the check that

$$\sum_{j \in N_h} \gamma_{ij} = 0, \tag{4.1}$$

namely, matrix  $\{\gamma_{ij}\}_{i,j \in N_h}$  satisfies “zero row-sum” condition. The same is true for the matrix  $\{\gamma_{ij}^*\}_{i,j \in N_h}$ . Suppose  $(y_{l,i}, v_{l,i})_{l=1}^M$  be the set of integration points

and corresponding weights of an  $M$ -point quadrature rule  $f_{\omega_i}^s$ . Then

$$\begin{aligned}
\sum_{j \in N_h} \gamma_{ij}^* &= \sum_{j \in N_h} \int_{\omega_i}^s A \nabla \phi_j \cdot \nabla \phi_i \, dx \\
&= \sum_{j \in N_h} \sum_{l=1}^M A(y_{l,i}) \nabla \phi_j(y_{l,i}) \cdot \nabla \phi_i(y_{l,i}) v_{l,i} \\
&= \sum_{l=1}^M A(y_{l,i}) \nabla \left[ \sum_{j \in N_h} \phi_j(y_{l,i}) \right] \cdot \nabla \phi_i(y_{l,i}) v_{l,i} \\
&= \sum_{l=1}^M A(y_{l,i}) \nabla 1 \cdot \nabla \phi_i(y_{l,i}) v_{l,i} = 0.
\end{aligned} \tag{4.2}$$

We note that (4.2) was an assumption on the quadrature rule in [5], where  $\gamma_{ij}^*$  was defined by using quadrature on  $\omega_i \cap \omega_j$ . This is one of the reasons that we defined  $\gamma_{ij}^*$  by numerically integrating over  $\omega_i$  in this paper (also in [6]) so that (4.2) is automatically satisfied.  $\blacksquare$

Let  $v_h = \sum_{i \in N_h} v_i \phi_i$  and  $w_h = \sum_{i \in N_h} w_i \phi_i$  be arbitrary elements in  $V_h$ . Then

$$\begin{aligned}
B_1(v_h, w_h) &= \sum_{i,j \in N_h} v_i \gamma_{ji} w_j, & B_0(v_h, w_h) &= \sum_{i,j \in N_h} v_i \sigma_{ji} w_j, \\
B(v_h, w_h) &= \sum_{i,j \in N_h} v_i (\gamma_{ji} + \sigma_{ji}) w_j, & \text{and } L(v_h) &= \sum_{i \in N_h} v_i l_i.
\end{aligned}$$

Therefore, we naturally define

$$B_1^*(v_h, w_h) \equiv \sum_{i,j \in N_h} v_i \gamma_{ji}^* w_j, \quad B_0^*(v_h, w_h) \equiv \sum_{i,j \in N_h} v_i \sigma_{ji}^* w_j, \tag{4.3}$$

$$B^*(v_h, w_h) \equiv \sum_{i,j \in N_h} v_i (\gamma_{ji}^* + \sigma_{ji}^*) w_j, \quad \text{and } L^*(v_h) \equiv \sum_{i \in N_h} v_i l_i^*. \tag{4.4}$$

From this definition, functional  $L^*(\cdot)$  is linear on  $V_h$  and the forms  $B_1^*(\cdot, \cdot)$ ,  $B_0^*(\cdot, \cdot)$ ,  $B^*(\cdot, \cdot)$  are bilinear on  $V_h \times V_h$ . Also from (4.2) and (4.1), it is clear that

$$B_1^*(1, \phi_i) = 0 = B_1(1, \phi_i), \quad \forall i \in N_h. \tag{4.5}$$

But it is important to note that the matrix  $\{\gamma_{ij}^*\}_{i,j \in N_h}$  may not be symmetric (in contrast to  $\{\gamma_{ij}\}_{i,j \in N_h}$ ), since

$$\gamma_{ij}^* = \int_{\omega_i}^s A \nabla \phi_j \cdot \nabla \phi_i \, dx \neq \int_{\omega_j}^s A \nabla \phi_i \cdot \nabla \phi_j \, dx = \gamma_{ji}^*.$$

Therefore,  $B_1^*(\phi_i, 1)$ ,  $\forall i \in N_h$  may not be zero. Similarly, we can show that the matrix  $\{\sigma_{ij}^*\}_{i,j \in N_h}$  may not be symmetric, and consequently, the matrix  $\{\gamma_{ij}^* + \sigma_{ij}^*\}_{i,j \in N_h}$  may not be symmetric.

The meshless method with numerical quadrature to approximate the solution of (2.3) is given by

$$\begin{aligned} &\text{Find } u_h^* \in V_h \text{ such that} \\ &B^*(u_h^*, v_h) = L^*(v_h), \quad \forall v_h \in V_h, \end{aligned} \quad (4.6)$$

where  $B^*(\cdot, \cdot)$  and  $L^*(\cdot)$  is defined in (4.4). We note that the bilinear form  $B^*(\cdot, \cdot)$  is not symmetric.

Next, we state certain assumptions on the quadrature used in the meshless method. Some of these assumptions were given in [6]. We include these assumptions also in this paper for completeness.

**QA 4.1** There exist positive constants  $\eta$  and  $\tau$ , small enough and independent of  $i$  and  $h$ , such that

$$\left| \int_{\omega_i}^t \varrho \, dx - \int_{\omega_i}^t \varrho \, dx \right| \leq \eta |\omega_i| \|\varrho\|_{L_\infty(\omega_i)}, \quad t = s, m, l, \quad (4.7)$$

and

$$\left| \int_{\partial\omega_i \cap \Gamma} \vartheta \, ds - \int_{\partial\omega_i \cap \Gamma} \vartheta \, ds \right| \leq \tau |\partial\omega_i \cap \Gamma| \|\vartheta\|_{L_\infty(\partial\omega_i \cap \Gamma)} \quad (4.8)$$

for a class of functions  $\varrho \in W^{m_1, \infty}(\omega_i)$  and  $\vartheta \in W^{m_2, \infty}(\partial\omega_i \cap \Gamma)$  satisfying

$$\|D^\alpha \varrho\|_{L_\infty(\omega_i)} \leq C(h_i)^{-|\alpha|} \|\varrho\|_{L_\infty(\omega_i)}, \quad |\alpha| \leq m_1 \quad (4.9)$$

and

$$\|D^\alpha \vartheta\|_{L_\infty(\partial\omega_i \cap \Gamma)} \leq C(h_i)^{-|\alpha|} \|\vartheta\|_{L_\infty(\partial\omega_i \cap \Gamma)}, \quad |\alpha| \leq m_2 \quad (4.10)$$

where  $C > 0$  is independent of  $i \in N_h$  and  $m_1, m_2 > 1$  may depend on the numerical integration rules and the assumption **A4** in Section 2.

**Remark 4.2** The constants  $\eta$  and  $\tau$  are associated with the numerical integration rules. It is possible to choose numerical integration rules (e.g., by taking more integration points) such that  $\eta$  and  $\tau$  are small enough. We refer to Remark 3.3 of [6] for specific examples. We mention that in all the numerically approximated integrals in this paper, the integrands satisfy the conditions (4.9) and (4.10). ■

**QA 4.2** For each  $i \in N_h$ , let  $G_i^* : \widetilde{C}^1(\bar{\omega}_i) \rightarrow \mathbb{R}$  be a linear functional given by

$$G_i^*(\tilde{v}) = \int_{\omega_i}^s \tilde{v} \cdot \nabla \phi_i \, dx + \int_{\omega_i}^l \nabla \cdot \tilde{v} \phi_i \, dx - \int_{\partial\omega_i \cap \Gamma} \tilde{v} \cdot \vec{n} \phi_i \, ds \quad (4.11)$$

where  $\vec{n}$  is the outward normal to  $\partial\omega_i \cap \Gamma$ . We assume that

$$G_i^*(\tilde{p}) = 0, \quad \forall \tilde{p} \in \widetilde{\mathcal{P}}^{k-1} \quad (4.12)$$

where  $\mathcal{P}^{k-1}$  is the space of polynomials of degree  $k-1$ .

**Remark 4.3** For each  $i \in N_h$ , we consider linear functional  $G_i : \widetilde{H}^1(\omega_i) \rightarrow \mathbb{R}$  given by

$$G_i(\tilde{v}) = \int_{\omega_i} \tilde{v} \cdot \nabla \phi_i \, dx + \int_{\omega_i} \nabla \cdot \tilde{v} \phi_i \, dx - \int_{\partial\omega_i \cap \Gamma} \tilde{v} \cdot \vec{n} \phi_i \, ds \quad (4.13)$$

It is clear from the Green Theorem that

$$G_i(\tilde{p}) = 0, \quad \forall \tilde{p} \in \widetilde{\mathcal{P}}^{k-1} \quad (4.14)$$

Hence, the assumption (4.12) mimics (4.14) and could be viewed as a discrete version of the Green Theorem on a particular class of functions  $\widetilde{\mathcal{P}}^{k-1}$ . We will show how to construct the quadrature rules satisfying (4.12) later. ■

**Remark 4.4** We note that the assumption **QA 4.2**, in particular (4.12), is a generalized version of a similar assumption QA3 used in [6] in the case when  $A = I$ . In fact, a numerical integration rule satisfying (4.12) also satisfies QA3 of [6], but not vice versa. Furthermore, we mention that for problems with non-constant coefficients  $A(x)$ , a direct use of the ideas presented in [6] will require the underlying numerical integration rule to satisfy a modified version of the assumption QA3 of [6] involving  $A(x)$ . Numerical integration rules, satisfying this modified assumption, will depend on  $A(x)$ , i.e., will be problem dependent. The assumption **QA 4.2** in this paper does not require the quadrature rules to depend on  $A(x)$ . ■

**QA 4.3** For each  $i \in N_h$ , we assume  $f_{\omega_i}^m = f_{\omega_i}^l$ , i.e., the elements of the mass matrix and the volume integrals in the elements the load vector are computed using the same integration rule.

**Remark 4.5** The integration rules  $f_{\omega_i}^s$  and  $f_{\omega_i}^l$  may be different; we do not need any other assumption on the boundary integration formulas  $f_{\partial\omega_i \cap \Gamma}$ .

**QA 4.4** There is a constant  $C > 0$  such that for  $\eta$  small enough,

$$|B_1^*(w_h, v_h)| \leq C \|w_h\|_{H^1(\Omega)} \|v_h\|_{H^1(\Omega)}, \quad \forall w_h, v_h \in V_h, \quad (4.15)$$

and

$$B_1^*(v_h, v_h) \geq C \|v_h\|_{H^1(\Omega)}^2, \quad \forall v_h \in V_h. \quad (4.16)$$

**Remark 4.6** For  $A = I$ , the identity matrix, the inequalities (4.15) and (4.16) were proved in Lemma 3.1 of [6] under the condition  $\eta \leq Ch$  with  $C$  small enough. The proof of (4.15) and (4.16) for the non-constant coefficient matrix  $A(x)$ , under the same conditions, could be proved following the same idea (of Lemma 3.1 of [6]) and we do not present it here. In our computation, however, we observed that the condition  $\eta \leq Ch$  is not necessary to obtain (4.15) and (4.16); only a small  $\eta$  was needed. ■

**Lemma 4.1** *Suppose the quadrature satisfy the assumptions **QA 4.1** and **QA 4.4**. Then for  $\eta$ , small enough, there are constants  $C_1$  and  $C_2$ , independent of  $h$ , such that*

$$|B^*(w_h, v_h)| \leq C_1 \|w_h\|_{H^1(\Omega)} \|v_h\|_{H^1(\Omega)} \quad \text{and} \quad B^*(v_h, v_h) \geq C_2 \|v_h\|_{H^1(\Omega)}^2$$

for any  $w_h, v_h \in V_h$ .

*Proof.* Let  $w_h = \sum_{i \in N_h} w_i \phi_i$  and  $v_h = \sum_{i \in N_h} v_i \phi_i$  be in  $V_h$ . We first estimate  $|B_0(w_h, v_h) - B_0^*(w_h, v_h)|$ . For any  $i \in N_h$ , using (4.7), the assumption **A1**, and (3.5), we have

$$\begin{aligned} |B_0(w_h, \phi_i) - B_0^*(w_h, \phi_i)| &= \left| \int_{\omega_i}^m c w_h \phi_i dx - \int_{\omega_i}^m c w_h \phi_i dx \right| \\ &\leq C \sum_{j \in S_i} |w_j| \left| \int_{\omega_i}^m c \phi_j \phi_i dx - \int_{\omega_i}^m c \phi_j \phi_i dx \right| \\ &\leq C \eta |\omega_i| \|c \phi_j \phi_i\|_{L^\infty(\omega_i)} \left( \sum_{j \in S_i} |w_j|^2 \right)^{\frac{1}{2}} \sqrt{\kappa} \\ &\leq C \eta h^d h^{-\frac{d}{2}} \|w_h\|_{L_2(\omega_i)} \sqrt{\kappa}. \end{aligned} \quad (4.17)$$

Therefore, squaring both sides of the above inequality and summing over all  $i \in N_h$ , we get

$$\begin{aligned} |B_0(w_h, v_h) - B_0^*(w_h, v_h)| &\leq \left( \sum_{i \in N_h} [B_0(w_h, \phi_i) - B_0^*(w_h, \phi_i)]^2 \right)^{\frac{1}{2}} \left( \sum_{i \in N_h} v_i^2 \right)^{\frac{1}{2}} \\ &\leq C \eta h^{\frac{d}{2}} \left( \sum_{i \in N_h} \|w_h\|_{L_2(\omega_i)}^2 \right)^{\frac{1}{2}} C h^{-\frac{d}{2}} \|v_h\|_{L_2(\Omega)} \\ &\leq C \eta \|w_h\|_{L_2(\Omega)} \|v_h\|_{L_2(\Omega)}, \end{aligned} \quad (4.18)$$

where the second and the last inequalities were obtained from (3.8) and the assumption **A1**, respectively.

Finally, from the assumption (4.4) and (4.18), we get

$$\begin{aligned} |B^*(w_h, v_h)| &\leq |B_1^*(w_h, v_h)| + |B_0(w_h, v_h)| + |B_0(w_h, v_h) - B_0^*(w_h, v_h)| \\ &\leq C \|w_h\|_{H^1(\Omega)} \|v_h\|_{H^1(\Omega)} + C(1 + \eta) \|w_h\|_{L_2(\Omega)} \|v_h\|_{L_2(\Omega)} \\ &\leq C(1 + \eta) \|w_h\|_{H^1(\Omega)} \|v_h\|_{H^1(\Omega)} \end{aligned}$$

and from (2.2)

$$\begin{aligned} |B^*(v_h, v_h)| &\geq B_1^*(v_h, v_h) + B_0(v_h, v_h) - |B_0(v_h, v_h) - B_0^*(v_h, v_h)| \\ &\geq C \|v_h\|_{H^1(\Omega)}^2 + \alpha \|v_h\|_{L_2(\Omega)}^2 - C \eta \|v_h\|_{L_2(\Omega)}^2 \\ &\geq \min\{C, \alpha - C\eta\} \|v_h\|_{H^1(\Omega)}^2. \end{aligned}$$

We get the desired result by considering  $\eta < \alpha/C$ .  $\square$

It is clear from Lemma 4.1 that the bilinear form  $B^*(\cdot, \cdot)$  is bounded and coercive, and therefore from the Lax-Milgram lemma we conclude that the problem (4.6) has a unique solution  $u_h^* \in V_h$ .

**Remark 4.7** It is instructive to illustrate the assumption **QA 4.2**, i.e., (4.12) in simpler situations. Let  $\Omega \subset \mathbb{R}^2$  and  $k = 1$ , then  $\tilde{\mathcal{P}}^{k-1} = \tilde{\mathcal{P}}^0 = \text{span}\{[1, 0], [0, 1]\}$ . Considering  $\tilde{p}(x_1, x_2) = [1, 0]$  in (4.12), we get

$$G_i^*([1, 0]) = \int_{\omega_i}^s \frac{\partial \phi_i}{\partial x_1} dx - \int_{\partial\omega_i \cap \Gamma} n_1 \phi_i ds = 0, \quad i \in N_h, \quad (4.19)$$

where  $\vec{n} = [n_1, n_2]$ . Similarly, considering  $\tilde{p}(x_1, x_2) = [0, 1]$  in (4.12), we get

$$G_i^*([0, 1]) = \int_{\omega_i}^s \frac{\partial \phi_i}{\partial x_2} dx - \int_{\partial\omega_i \cap \Gamma} n_2 \phi_i ds = 0, \quad i \in N_h \quad (4.20)$$

Thus for  $k = 1$ , the quadrature must satisfy the two conditions (4.19) and (4.20) for each  $i \in N_h$ . In particular, the quadrature must satisfy

$$\int_{\omega_i}^s \nabla \phi_i dx = 0, \quad \forall i \in N_h''. \quad (4.21)$$

We now illustrate (4.12) for  $k = 2$ . In this case, we know that

$$\tilde{\mathcal{P}}^{k-1} = \tilde{\mathcal{P}}^1 = \text{span}\{[1, 0], [0, 1], [x_1, 0], [x_2, 0], [0, x_1], [0, x_2]\}.$$

Considering  $\tilde{p}(x_1, x_2) = [x_1, 0]$  in (4.12), we get

$$G_i([x_1, 0]) = \int_{\omega_i}^s x_1 \frac{\partial \phi_i}{\partial x_1} dx + \int_{\omega_i}^l \phi_i dx - \int_{\partial\omega_i \cap \Gamma} x_1 n_1 \phi_i ds = 0, \quad \forall i \in N_h \quad (4.22)$$

Similarly, considering  $\tilde{p}(x_1, x_2) = [x_2, 0]$ ,  $p(x_1, x_2) = [0, x_1]$ , and  $\tilde{p}(x_1, x_2) = [0, x_2]$  in (4.12), we get

$$G_i([x_2, 0]) = \int_{\omega_i}^s x_2 \frac{\partial \phi_i}{\partial x_1} dx - \int_{\partial\omega_i \cap \Gamma} x_2 n_1 \phi_i ds = 0, \quad \forall i \in N_h, \quad (4.23)$$

$$G_i([0, x_1]) = \int_{\omega_i}^s x_1 \frac{\partial \phi_i}{\partial x_2} dx - \int_{\partial\omega_i \cap \Gamma} x_1 n_2 \phi_i ds = 0, \quad \forall i \in N_h, \quad (4.24)$$

and

$$G_i([0, x_2]) = \int_{\omega_i}^s x_2 \frac{\partial \phi_i}{\partial x_2} dx + \int_{\omega_i}^l \phi_i dx - \int_{\partial\omega_i \cap \Gamma} x_2 n_2 \phi_i ds = 0, \quad \forall i \in N_h \quad (4.25)$$

Thus, for  $k = 2$ , the quadrature must satisfy (4.22)–(4.25) in addition to the assumptions (4.19) and (4.20).  $\blacksquare$

**Remark 4.8** It is clear from (4.19) and (4.20) that for  $k = 1$ , only the quadrature rule to compute the elements of the stiffness matrix, i.e.,  $f_{\omega_i}^s$ , has to satisfy the assumption **QA 4.2**; the quadrature rule to compute the elements of the mass matrix and the volume integrals in the elements of the load vector, i.e.,  $f_{\omega_i}^m$

(it is same as  $f_{\omega_i}^l$ ), do not have to satisfy **QA 4.2** and it could be any accurate rule satisfying assumption **QA 4.1**. For  $k = 2$ , the conditions (4.22) and (4.25) indicate that  $f_{\omega_i}^s$  and  $f_{\omega_i}^l$  must be related. Even in this situation,  $f_{\omega_i}^l$  (same as  $f_{\omega_i}^m$ ) could be any accurate quadrature rule, but  $f_{\omega_i}^s$  has to be satisfy (4.22) and (4.25). We will obtain  $f_{\omega_i}^s$  later in the paper with this feature.  $\blacksquare$

## 5 Effect of numerical integration

In this section, we will investigate the effect of numerical integration on the meshless method; in particular, we will estimate the error  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u$  is the solution of the problem (2.3) and  $u_h^*$  is the solution of the meshless method (4.6) with numerical integration. We recall from Theorem 3.2 that  $\|u - u_h\|_{H^1(\Omega)} = \mathcal{O}(h^k)$ , where  $u_h$  is the solution of (3.11) — the meshless method with exact integration. We will show in this section that  $\|u - u_h^*\|_{H^1(\Omega)} \neq \mathcal{O}(h^k)$  in general, and the error depends on the quadrature parameters  $\eta$  and  $\tau$ , defined in (4.7) and (4.8), respectively. We will assume in this section that the exact solution  $u$  of (2.3) is smooth, i.e.,  $u \in C^{k+1}(\bar{\Omega})$ ; this will enable us to focus only on numerical integration and will allow us to present the main ideas effectively.

It is well-known that Strang's Lemma ([27, 15]) is one of the main tools to study the perturbation in the solution of a Galerkin method due to variational crimes, e.g., numerical integration in a Galerkin method. We present a slight variation of the Strang's Lemma in the following result, which will provide us with an abstract framework to study the error  $u - u_h^*$ .

**Lemma 5.1** *Suppose the quadrature rules satisfy the conditions in the lemma 4.1, and  $u$  and  $u_h^*$  are the solutions of the variational problems (2.3) and (4.6), respectively. Then there is a constant  $C > 0$ , independent of  $h$ , such that, for any  $w_h \in V_h$ ,*

$$\begin{aligned} \|u - u_h^*\|_{H^1(\Omega)} &\leq C \|u - w_h\|_{H^1(\Omega)} \\ &+ \sup_{v_h \in V_h} \frac{\left| [B(w_h, v_h) - L(v_h)] - [B^*(w_h, v_h) - L^*(v_h)] \right|}{\|v_h\|_{H^1(\Omega)}}. \end{aligned}$$

*Proof.* Let  $w_h \in V_h$  be arbitrary. Using the coercivity of the bilinear form  $B^*(\cdot, \cdot)$  (see Lemma 4.1), we have

$$\begin{aligned} C \|u_h^* - w_h\|_{H^1(\Omega)}^2 &\leq B^*(u_h^* - w_h, u_h^* - w_h) \\ &= B(u - w_h, u_h^* - w_h) \\ &\quad + B(w_h, u_h^* - w_h) - B^*(w_h, u_h^* - w_h) \\ &\quad - B(u, u_h^* - w_h) + B^*(u_h^*, u_h^* - w_h) \\ &= B(u - w_h, u_h^* - w_h) \\ &\quad + [B(w_h, u_h^* - w_h) - L(u_h^* - w_h)] \\ &\quad - [B^*(w_h, u_h^* - w_h) - L^*(u_h^* - w_h)]. \end{aligned}$$



Therefore, dividing the above inequality by  $\|u_h^* - w_h\|_{H^1(\Omega)}$  and using the boundedness of  $B(\cdot, \cdot)$ , we get

$$\begin{aligned} \|u_h^* - w_h\|_{H^1(\Omega)} &\leq C\|u - w_h\|_{H^1(\Omega)} \\ &\quad + \sup_{v_h \in V_h} \frac{\left| [B(w_h, v_h) - L(v_h)] - [B^*(w_h, v_h) - L^*(v_h)] \right|}{\|v_h\|_{H^1(\Omega)}} \end{aligned}$$

Now, using the triangle inequality, we immediately get

$$\begin{aligned} \|u - u_h^*\|_{H^1(\Omega)} &\leq (C + 1)\|u - w_h\|_{H^1(\Omega)} \\ &\quad + \sup_{v_h \in V_h} \frac{\left| [B(w_h, v_h) - L(v_h)] - [B^*(w_h, v_h) - L^*(v_h)] \right|}{\|v_h\|_{H^1(\Omega)}}, \end{aligned}$$

which is the desired result.  $\square$

**Remark 5.1** It is clear from lemma 5.1 that we need to estimate the consistency errors

$$\sup_{v_h \in V_h} \frac{\left| [B(w_h, v_h) - L(v_h)] - [B^*(w_h, v_h) - L^*(v_h)] \right|}{\|v_h\|_{H^1(\Omega)}} \quad (5.1)$$

to estimate the error  $\|u - u_h^*\|_{H^1(\Omega)}$ . We note that in the Strang's Lemma as presented in [15], this term is further divided into two terms

$$\sup_{v_h \in V_h} \frac{\left| B(w_h, v_h) - B^*(w_h, v_h) \right|}{\|v_h\|_{H^1(\Omega)}} \quad \text{and} \quad \sup_{v_h \in V_h} \frac{\left| L(v_h) - L^*(v_h) \right|}{\|v_h\|_{H^1(\Omega)}}.$$

Keeping the terms together, as in (5.1), is crucial for our analysis of the effect of numerical integration in the meshless method.  $\blacksquare$

We now present some notions and associated results that we will use later in this section. We first define a norm and semi-norm of the matrix  $A$ ; recall that we assumed  $a_{ij}(x) \in C^k(\bar{\Omega})$ ,  $\forall i, j = 1, 2, \dots, d$ . Suppose  $D \subset \Omega$  be a domain and let

$$|A|_{W^{l, \infty}(D)} \equiv \max \left\{ \sum_{j=1}^d |a_{ij}|_{W^{l, \infty}(D)} : 1 \leq i \leq d \right\}$$

$$\text{and} \quad \|A\|_{W^{l, \infty}(D)} \equiv \max \{ |A|_{W^{m, \infty}(D)} : 0 \leq m \leq l \},$$

for any non-negative integer  $l \leq k$ .

**Lemma 5.2** For  $0 \leq l \leq k$ , there exists a constant  $C > 0$ , depending only on  $l$  and  $d$ , such that

$$\|A \tilde{v}\|_{W^{l, \infty}(D)} \leq C \|A\|_{W^{l, \infty}(D)} \|\tilde{v}\|_{W^{l, \infty}(D)}, \quad \forall \tilde{v} \in \widetilde{W}^{k, \infty}(D). \quad (5.2)$$

*Proof.* Let  $0 \leq \ell \leq l$  and suppose  $\tilde{v} = [v_j]_{j=1}^d$ . Then using Leibnitz formula, we have

$$\begin{aligned}
|A\tilde{v}|_{W^{\ell,\infty}(D)} &= \max_{1 \leq i \leq d} \left\{ \left| \sum_{j=1}^d a_{ij} v_j \right|_{W^{\ell,\infty}(D)} \right\} \\
&\leq C \max_{1 \leq i \leq d} \left\{ \sum_{j=1}^d \sum_{m=0}^{\ell} |a_{ij}|_{W^{\ell-m,\infty}(D)} |v_j|_{W^{m,\infty}(D)} \right\} \\
&\leq C \sum_{m=0}^{\ell} |\tilde{v}|_{W^{m,\infty}(D)} \max_{1 \leq i \leq d} \left\{ \sum_{j=1}^d |a_{ij}|_{W^{\ell-m,\infty}(D)} \right\} \\
&= C \sum_{m=0}^{\ell} |\tilde{v}|_{W^{m,\infty}(D)} |A|_{W^{\ell-m,\infty}(D)} \\
&\leq C \|A\|_{W^{\ell,\infty}(D)} \|\tilde{v}\|_{W^{\ell,\infty}(D)} \leq C \|A\|_{W^l,\infty(D)} \|\tilde{v}\|_{W^l,\infty(D)},
\end{aligned}$$

where the constant  $C$  only depends on  $l$  and  $d$ . Therefore,

$$\|A\tilde{v}\|_{W^l,\infty(D)} \leq C \|A\|_{W^l,\infty(D)} \|\tilde{v}\|_{W^l,\infty(D)},$$

which is the desired result.  $\square$

We now present the next result. For a smooth function  $v$  and  $i \in N_h$ , let

$$T_i^{k-1}v(x) = \sum_{|\alpha| \leq k-1} \frac{D^\alpha v(\bar{x}_i)}{\alpha!} (x - \bar{x}_i)^\alpha$$

be the  $(k-1)^{th}$  degree Taylor polynomial of  $v$  associated with the center  $\bar{x}_i$  of the ball  $o_i \subset \omega_i$  (recall in Section 2 that  $\omega_i$  is star-shaped with respect to the ball  $o_i$ ). It is well known that ([10])

$$|v - T_i^{k-1}v|_{W^j,\infty(\omega_i)} \leq \frac{Ch^{k-j}}{(k-j)!} \|v\|_{W^{k,\infty}(\omega_i)}, \quad j = 0, 1, \dots, k \quad (5.3)$$

For a smooth vector-valued function  $\tilde{v} = [v_j]_{j=1}^d$  we define

$$\tilde{T}_i^{k-1}\tilde{v}(x) = [T_i^{k-1}v_j]_{j=1}^d.$$

$\tilde{T}_i^{k-1}\tilde{v}(x)$  is also vector-valued function with its components being the  $(k-1)^{th}$  degree Taylor polynomials of the corresponding components of  $\tilde{v}$ , centered at  $\bar{x}_i$ . We will refer to  $\tilde{T}_i^{k-1}\tilde{v}(x)$  as the Taylor polynomial of  $\tilde{v}$  associated with  $\bar{x}_i$ .

We define

$$\tilde{R}_i \equiv A\nabla\mathcal{I}_h u - \tilde{T}_i^{k-1}(A\nabla\mathcal{I}_h u), \quad (5.4)$$

where  $\mathcal{I}_h u$  is the  $V_h$ -interpolant of  $u$ , defined in (3.12) through (3.13). Clearly  $\tilde{R}_i$  is the ‘‘remainder’’ of the Taylor polynomial  $\tilde{T}_i^{k-1}\tilde{v}(x)$  with  $\tilde{v} = A\nabla\mathcal{I}_h u$ .

**Lemma 5.3** *Let  $0 \leq j \leq k$ . Then there exists a constant  $C > 0$  such that*

$$|\tilde{R}_i|_{W^{j,\infty}(\omega_i)} \leq Ch^{k-j} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)} \quad (5.5)$$

*Proof.* Let  $\tilde{v} \in C^k(\bar{\omega}_i)$ . Then from the definition of norm of vector valued functions and from (5.3), we immediately get

$$|\tilde{v} - \tilde{T}_i^{k-1}\tilde{v}|_{W^{j,\infty}(\omega_i)} \leq Ch^{k-j} \|\tilde{v}\|_{W^{k,\infty}(\omega_i)}, \quad j = 0, 1, \dots, k$$

Now substituting  $\tilde{v} = A\nabla\mathcal{I}_h u$  in the above inequality, and using (5.2) and (3.19), we get, for  $j = 0, 1, \dots, k$ ,

$$\begin{aligned} |\tilde{R}_i|_{W^{j,\infty}(\omega_i)} &\leq Ch^{k-j} \|A\nabla\mathcal{I}_h u\|_{W^{k,\infty}(\omega_i)} \\ &\leq Ch^{k-j} \|A\|_{W^{k,\infty}(\omega_i)} \|\nabla\mathcal{I}_h u\|_{W^{k,\infty}(\omega_i)} \\ &\leq Ch^{k-j} \|A\|_{W^{k,\infty}(\Omega)} \|\nabla\mathcal{I}_h u\|_{W^{k,\infty}(\Omega)} \\ &\leq Ch^{k-j} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)}, \end{aligned}$$

which is the desired result.  $\square$

The next lemma provides us with an estimate of the error in the numerical integration for a particular integrand, and it is an important ingredient in the proof of the main result of the paper. This result is a generalization of Lemma 4.2 in [6] in the context of variable coefficients  $A(x) = [a_{ij}(x)]_{1 \leq i, j \leq d}$ ; we mention that the matrix  $A(x) = I$  was considered in [6].

**Lemma 5.4** *For any  $i \in N_h$ , let  $G_i(\cdot)$  and  $G_i^*(\cdot)$  be functionals defined by (4.13), (4.11), respectively. Assume that the quadrature formulas satisfy the assumptions (4.7), (4.8), and (4.12). Then there exists a positive constant  $C$ , independent of  $h$  and  $i$ , such that, for  $i \in N_h''$ ,*

$$|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \leq C\eta h^{k+d-1} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)},$$

and, for  $i \in N_h'$ ,

$$|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \leq C(\eta + \tau) h^{k+d-1} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)}.$$

*Proof.* For  $i \in N_h$ , let  $\bar{x}_i$  be the center of the ball  $o_i \subset \omega_i$ . We expand the vector-valued function  $A\nabla\mathcal{I}_h u$  with respect to  $\bar{x}_i$  using (5.4) as

$$A\nabla\mathcal{I}_h u = \tilde{T}_i^{k-1}(A\nabla\mathcal{I}_h u) + \tilde{R}_i.$$

We note that  $\tilde{T}_i^{k-1}(A\nabla\mathcal{I}_h u) \in \tilde{P}^{k-1}$ . Therefore from the assumption on the quadrature (4.12) and the fact (4.14), we have

$$G_i^*(\tilde{T}_i^{k-1}(A\nabla\mathcal{I}_h u)) = 0 \quad \text{and} \quad G_i(\tilde{T}_i^{k-1}(A\nabla\mathcal{I}_h u)) = 0.$$

Hence, for  $i \in N'_h$ ,

$$\begin{aligned}
|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| &= |G_i(\tilde{R}_i) - G_i^*(\tilde{R}_i)| \\
&\leq \left| \int_{\omega_i} \tilde{R}_i \cdot \nabla \phi_i \, dx - \int_{\omega_i}^s \tilde{R}_i \cdot \nabla \phi_i \, dx \right| \\
&\quad + \left| \int_{\omega_i} \nabla \cdot \tilde{R}_i \phi_i \, dx - \int_{\omega_i}^l \nabla \cdot \tilde{R}_i \phi_i \, dx \right| \\
&\quad + \left| \int_{\partial\omega_i \cap \Gamma} \tilde{R}_i \cdot \tilde{n} \phi_i \, ds - \int_{\partial\omega_i \cap \Gamma} \tilde{R}_i \cdot \tilde{n} \phi_i \, ds \right|. \quad (5.6)
\end{aligned}$$

Also for  $i \in N''_h$ , recalling that  $\phi_i = 0$  on  $\partial\omega_i$ , we get

$$\begin{aligned}
|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| &= |G_i(\tilde{R}_i) - G_i^*(\tilde{R}_i)| \\
&\leq \left| \int_{\omega_i} \tilde{R}_i \cdot \nabla \phi_i \, dx - \int_{\omega_i}^s \tilde{R}_i \cdot \nabla \phi_i \, dx \right| \\
&\quad + \left| \int_{\omega_i} \nabla \cdot \tilde{R}_i \phi_i \, dx - \int_{\omega_i}^l \nabla \cdot \tilde{R}_i \phi_i \, dx \right|. \quad (5.7)
\end{aligned}$$

Now, from (5.6), the assumptions **QA 4.1**, **A4**, **A2**, and the remainder estimate (5.5), we obtain for  $i \in N'_h$

$$\begin{aligned}
|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| &= \eta |\tilde{R}_i \cdot \nabla \phi_i|_{L_\infty(\omega_i)} |\omega_i| + \eta |\nabla \cdot \tilde{R}_i \phi_i|_{L_\infty(\omega_i)} |\omega_i| \\
&\quad + \tau |\tilde{R}_i \cdot \tilde{n} \phi_i|_{L_\infty(\partial\omega_i \cap \Gamma)} |\partial\omega_i \cap \Gamma| \\
&\leq 2C\eta h^{k-1+d} \|A\|_{W^{k,\infty}(\omega_i)} \|u\|_{W^{k+1,\infty}(\omega_i)} \\
&\quad + C\tau h^{k-1+d} \|A\|_{W^{k,\infty}(\omega_i)} \|u\|_{W^{k+1,\infty}(\omega_i)} \\
&\leq C(\eta + \tau) h^{k-1+d} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)},
\end{aligned}$$

which is the desired result for  $i \in N'_h$ . Also using (5.6) and similar arguments as above, we get for  $i \in N''_h$

$$|G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \leq C\eta h^{k-1+d} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)},$$

which completes the proof.  $\square$

Now, we present our main result, where we estimate the energy norm of the error  $u - u_h^*$ ; recall that  $u_h^*$  is the unique solution of the meshless method (4.6) with numerical integration.

**Theorem 5.5** *Let  $u \in C^{k+1}(\bar{\Omega})$ ,  $a_{ij} \in C^k(\bar{\Omega})$ , for  $i, j = 1, 2, \dots, d$  and  $c \in C(\bar{\Omega})$ . Suppose the subspace  $V_h$  satisfies assumptions **A1-A5** and the quadrature schemes satisfy **QA1-QA4**. Then, for  $\eta$  small enough, there is a positive constant  $C$ , independent of  $u$ ,  $\eta$ ,  $\tau$ , and  $h$ , such that*

$$\begin{aligned}
\|u - u_h^*\|_{H^1(\Omega)} &\leq Ch^k \|u\|_{W^{k+1,\infty}(\Omega)} + \left[ C\eta (\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L_\infty(\Omega)} h^2) \right. \\
&\quad \left. + (\eta + \tau) (\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L_\infty(\Omega)} h^2) h \right] h^{k-1} \|u\|_{W^{k+1,\infty}(\Omega)}. \quad (5.8)
\end{aligned}$$

*Proof.* First, we substitute  $w_h = \mathcal{I}_h u$  in the result of Lemma 5.1 and get

$$\begin{aligned} \|u - u_h^*\|_{H^1(\Omega)} &\leq C \|u - \mathcal{I}_h u\|_{H^1(\Omega)} \\ &+ \sup_{v_h \in V_h} \frac{\left| [B(\mathcal{I}_h u, v_h) - L(v_h)] - [B^*(\mathcal{I}_h u, v_h) - L^*(v_h)] \right|}{\|v_h\|_{H^1(\Omega)}}. \end{aligned} \quad (5.9)$$

We will now estimate the second part of the RHS of (5.9) to prove (5.8). For any  $v_h = \sum_{i \in N_h} v_i \phi_i \in V_h$ , we have

$$\begin{aligned} &[B(\mathcal{I}_h u, v_h) - L(v_h)] - [B^*(\mathcal{I}_h u, v_h) - L^*(v_h)] \\ &= \sum_{i \in N_h} v_i ([B(\mathcal{I}_h u, \phi_i) - L(\phi_i)] - [B^*(\mathcal{I}_h u, \phi_i) - L^*(\phi_i)]). \end{aligned} \quad (5.10)$$

For simplicity, in the rest of the proof we denote

$$E_i \equiv \left| [B(\mathcal{I}_h u, \phi_i) - L(\phi_i)] - [B^*(\mathcal{I}_h u, \phi_i) - L^*(\phi_i)] \right|.$$

Therefore, from (5.10), (3.9), (3.8), and (3.10), we get

$$\begin{aligned} &\left| [B(\mathcal{I}_h u, v_h) - L(v_h)] - [B^*(\mathcal{I}_h u, v_h) - L^*(v_h)] \right| \\ &\leq \sup_{i \in N'_h} E_i \sum_{i \in N'_h} |v_i| + \sup_{i \in N''_h} E_i \sum_{i \in N''_h} |v_i| \\ &\leq C \sup_{i \in N'_h} E_i \left( \sum_{i \in N'_h} |v_i|^2 \right)^{\frac{1}{2}} |N'_h|^{\frac{1}{2}} + C \sup_{i \in N''_h} E_i \left( \sum_{i \in N''_h} |v_i|^2 \right)^{\frac{1}{2}} |N''_h|^{\frac{1}{2}} \\ &\leq C \sup_{i \in N'_h} E_i h^{-(d-1)} \|v_h\|_{L_2(\Gamma)} + C \sup_{i \in N''_h} E_i h^{-d} \|v_h\|_{L_2(\Omega)} \\ &\leq C \sup_{i \in N'_h} E_i h^{-(d-1)} \|v_h\|_{H^1(\Omega)} + C \sup_{i \in N''_h} E_i h^{-d} \|v_h\|_{L_2(\Omega)} \end{aligned} \quad (5.11)$$

where the last inequality was obtained using the Trace theorem (see [10]). We will now estimate the terms  $E_i$ ,  $\forall i \in N_h$ . For any  $i \in N_h$ , we have from the problem (2.1)

$$\begin{aligned} \int_{\omega_i} f \phi_i dx &= - \int_{\omega_i} \nabla \cdot (A \nabla u) \phi_i dx + \int_{\omega_i} c u \phi_i dx \\ &= - \int_{\omega_i} \nabla \cdot (A \nabla \mathcal{I}_h u) \phi_i dx - \int_{\omega_i} \nabla \cdot [A \nabla (u - \mathcal{I}_h u)] \phi_i dx + \int_{\omega_i} c u \phi_i dx \end{aligned}$$

and

$$\begin{aligned} \int_{\partial \omega_i \cap \Gamma} g \phi_i ds &= \int_{\partial \omega_i \cap \Gamma} A \nabla u \cdot \vec{n} \phi_i ds \\ &= \int_{\partial \omega_i \cap \Gamma} A \nabla \mathcal{I}_h u \cdot \vec{n} \phi_i ds + \int_{\partial \omega_i \cap \Gamma} A \nabla (u - \mathcal{I}_h u) \cdot \vec{n} \phi_i ds. \end{aligned}$$

Now setting  $e_I \equiv u - \mathcal{I}_h u$  and recalling the definition (4.13) of the functional  $G_i$ , we get

$$\begin{aligned}
B(\mathcal{I}_h u, \phi_i) - L(\phi_i) &= B(\mathcal{I}_h u, \phi_i) - \int_{\omega_i} f \phi_i dx - \int_{\partial\omega_i \cap \Gamma} g \phi_i ds \\
&= \int_{\omega_i} A \nabla \mathcal{I}_h u \cdot \nabla \phi_i dx + \int_{\omega_i} \nabla \cdot (A \nabla \mathcal{I}_h u) \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla \mathcal{I}_h u \cdot \vec{n} \phi_i ds \\
&\quad + \int_{\omega_i} \nabla \cdot [A \nabla (u - \mathcal{I}_h u)] \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla (u - \mathcal{I}_h u) \cdot \vec{n} \phi_i ds \\
&\quad + \int_{\omega_i} c \mathcal{I}_h u \phi_i dx - \int_{\omega_i} c u \phi_i dx \\
&= G_i(A \nabla \mathcal{I}_h u) + \int_{\omega_i} \nabla \cdot (A \nabla e_I) \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla e_I \cdot \vec{n} \phi_i ds \\
&\quad - \int_{\omega_i} c e_I \phi_i dx. \tag{5.12}
\end{aligned}$$

Similarly, again from the problem (2.1) and the quadratures developed in Section 4, we have

$$\begin{aligned}
\int_{\omega_i}^l f \phi_i dx &= - \int_{\omega_i}^l \nabla \cdot (A \nabla u) \phi_i dx + \int_{\omega_i}^l c u \phi_i dx \\
&= - \int_{\omega_i}^l \nabla \cdot (A \nabla \mathcal{I}_h u) \phi_i dx - \int_{\omega_i}^l \nabla \cdot [A \nabla (u - \mathcal{I}_h u)] \phi_i dx + \int_{\omega_i}^l c u \phi_i dx
\end{aligned}$$

and

$$\begin{aligned}
\int_{\partial\omega_i \cap \Gamma} g \phi_i ds &= \int_{\partial\omega_i \cap \Gamma} A \nabla u \cdot \vec{n} \phi_i ds \\
&= \int_{\partial\omega_i \cap \Gamma} A \nabla \mathcal{I}_h u \cdot \vec{n} \phi_i ds + \int_{\partial\omega_i \cap \Gamma} A \nabla (u - \mathcal{I}_h u) \cdot \vec{n} \phi_i ds.
\end{aligned}$$

Then recalling the definition (4.11) of the functional  $G_i^*$ , we get

$$\begin{aligned}
B^*(\mathcal{I}_h u, \phi_i) - L^*(\phi_i) &= B^*(\mathcal{I}_h u, \phi_i) - \int_{\omega_i}^l f \phi_i dx - \int_{\partial\omega_i \cap \Gamma} g \phi_i ds \\
&= \int_{\omega_i}^s A \nabla \mathcal{I}_h u \cdot \nabla \phi_i dx + \int_{\omega_i}^l \nabla \cdot (A \nabla \mathcal{I}_h u) \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla \mathcal{I}_h u \cdot \vec{n} \phi_i ds \\
&\quad + \int_{\omega_i}^l \nabla \cdot [A \nabla (u - \mathcal{I}_h u)] \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla (u - \mathcal{I}_h u) \cdot \vec{n} \phi_i ds \\
&\quad + \int_{\omega_i}^m c \mathcal{I}_h u \phi_i dx - \int_{\omega_i}^l c u \phi_i dx \\
&= G_i^*(A \nabla \mathcal{I}_h u) + \int_{\omega_i}^l \nabla \cdot (A \nabla e_I) \phi_i dx - \int_{\partial\omega_i \cap \Gamma} A \nabla e_I \cdot \vec{n} \phi_i ds \\
&\quad - \int_{\omega_i}^l c e_I \phi_i dx, \tag{5.13}
\end{aligned}$$

where the last equality is due to the assumption  $f_{\omega_i}^l = f_{\omega_i}^m$ . Therefore, from (5.12), (5.13) and the assumptions (4.7), (4.8), we get the following estimates for  $i \in N'_h$ , namely,

$$\begin{aligned}
E_i &\leq |G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \\
&\quad + \left| \int_{\omega_i} \nabla \cdot (A\nabla e_I)\phi_i dx - \int_{\omega_i}^l \nabla \cdot (A\nabla e_I)\phi_i dx \right| \\
&\quad + \left| \int_{\partial\omega_i \cap \Gamma} A\nabla e_I \cdot \vec{n}\phi_i ds - \int_{\partial\omega_i \cap \Gamma}^l A\nabla e_I \cdot \vec{n}\phi_i ds \right| \\
&\quad + \left| \int_{\omega_i} c e_I \phi_i dx - \int_{\omega_i}^l c e_I \phi_i dx \right| \\
&\leq |G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| + \eta|\omega_i| |(A\nabla e_I)\phi_i|_{W^{1,\infty}(\omega_i)} \\
&\quad + \tau|\partial\omega_i \cap \Gamma| \|A\nabla e_I \cdot \vec{n}\phi_i\|_{L^\infty(\partial\omega_i \cap \Gamma)} + \eta|\omega_i| \|c e_I \phi_i\|_{L^\infty(\omega_i)}, \quad (5.14)
\end{aligned}$$

and similarly, for  $i \in N''_h$ , recalling that  $\phi_i = 0$  on  $\partial\omega_i$ , we have

$$\begin{aligned}
E_i &\leq |G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \\
&\quad + \eta|\omega_i| |(A\nabla e_I)\phi_i|_{W^{1,\infty}(\omega_i)} + \eta|\omega_i| \|c e_I \phi_i\|_{L^\infty(\omega_i)}, \quad (5.15)
\end{aligned}$$

Now, from (5.2), the interpolation error (3.14), and the boundedness of  $\phi_i$ , it immediately follows that for  $i \in N_h$ ,

$$\begin{aligned}
|(A\nabla e_I)\phi_i|_{W^{1,\infty}(\omega_i)} &\leq C \|A\|_{W^{1,\infty}(\omega_i)} h^{k-1} \|u\|_{W^{k+1,\infty}(\Omega)} \\
\text{and } \|c e_I \phi_i\|_{L^\infty(\omega_i)} &\leq C \|c\|_{L^\infty(\omega_i)} h^{k+1} \|u\|_{W^{k+1,\infty}(\Omega)},
\end{aligned}$$

and for  $i \in N'_h$ ,

$$\|A\nabla e_I \cdot \vec{n}\phi_i\|_{L^\infty(\partial\omega_i \cap \Gamma)} \leq C \|A\|_{L^\infty(\omega_i)} h^k \|u\|_{W^{k+1,\infty}(\Omega)}.$$

Therefore, from (5.14), (5.15), the assumption **A2**, and Lemma 5.4, we get

$$E_i \leq \begin{cases} C[(\eta + \tau)(\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L^\infty(\Omega)} h^2)] h^{k+d-1} \|u\|_{W^{k+1,\infty}(\Omega)}, & i \in N'_h; \\ C[\eta(\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L^\infty(\Omega)} h^2)] h^{k+d-1} \|u\|_{W^{k+1,\infty}(\Omega)}, & i \in N''_h. \end{cases} \quad (5.16)$$

Finally, from (5.9), the interpolation error (3.14), (5.11), and (5.16), we get

$$\begin{aligned}
\|u - u_h^*\|_{H^1(\Omega)} &\leq C h^k \|u\|_{W^{k+1,\infty}(\Omega)} + \left[ C\eta(\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L^\infty(\Omega)} h^2) \right. \\
&\quad \left. + (\eta + \tau)(\|A\|_{W^{k,\infty}(\Omega)} + \|c\|_{L^\infty(\Omega)} h^2) h \right] h^{k-1} \|u\|_{W^{k+1,\infty}(\Omega)},
\end{aligned}$$

which is the required result.  $\square$

**Remark 5.2** The result (5.8) of Theorem 5.5 shows that  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}[h^k + (\eta + \tau)h^k + \eta h^{k-1}]$ . Thus we do not have the optimal order of convergence

(compare with (3.18)). But if we consider numerical integration such that  $\eta = \mathcal{O}(h)$ , i.e, we use more accurate integration scheme as we refine  $h$ , we get back the optimal order of convergence  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h^k)$ . This feature of the meshless method is very different from the standard FEM, where the same numerical integration can be used for all values of  $h$  to obtain the optimal order of convergence. We further note that (5.8) indicates that for larger values of  $h$  (i.e., in the pre-asymptotic range), the error  $\|u - u_h^*\|_{H^1(\Omega)}$  may behave like  $\mathcal{O}(h^k)$ . But as  $h$  becomes smaller, we get  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h^{k-1})$ . We will show this feature in our numerical experiments.  $\blacksquare$

**Corollary 5.6** *Suppose all the assumptions in Theorem 5.5 hold, except for the assumption **QA 4.2**, which is replaced as follows: For a non-negative integer  $l < k$ ,*

$$G_i^*(\tilde{p}) = 0, \quad \forall \tilde{p} \in \tilde{\mathcal{P}}^{l-1} \text{ and } \forall i \in N_h; \quad (5.17)$$

for the case  $l = 0$ , we assume that the condition (5.17) is vacuous, namely, numerical integration rules satisfy only **QA 4.1**, **QA 4.3**, and **QA 4.4**. Then, for  $\eta$  small enough, there is a positive constant  $C$ , independent of  $u$ ,  $\eta$ ,  $\tau$ , and  $h$ , such that

$$\|u - u_h^*\|_{H^1(\Omega)} \leq C[h^k + (\eta + \tau)h^l + \eta h^{l-1}] \|u\|_{W^{k+1,\infty}(\Omega)}.$$

*Proof.* (Only a sketch) It can be easily shown by following the proof of the Lemma 5.4 that for  $0 \leq l \leq k$ ,

$$\begin{aligned} & |G_i(A\nabla\mathcal{I}_h u) - G_i^*(A\nabla\mathcal{I}_h u)| \\ & \leq \begin{cases} C(\eta + \tau)h^{l+d-1} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)}, & i \in N_h' \\ C\eta h^{l+d-1} \|A\|_{W^{k,\infty}(\Omega)} \|u\|_{W^{k+1,\infty}(\Omega)}, & i \in N_h'' \end{cases}. \end{aligned} \quad (5.18)$$

Moreover, for  $l = 0$ , we do not need to use the Taylor polynomial of  $A\nabla\mathcal{I}_h u$  (as in the proof of Lemma 5.4) to get, (5.18). Now, instead of using the result of Lemma 5.4, we use (5.18) in the proof of Theorem 5.5 to get the desired result.  $\square$

**Remark 5.3** The result in Corollary 5.6 shows that if the quadrature rules satisfy (4.12) of the assumption **QA 4.2** with  $k$  replaced by  $l$  and  $l < k$ , then  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h^{l-1})$ . Also, if the quadrature rules do not satisfy (4.12) (i.e.,  $l = 0$ ), then  $\|u - u_h^*\|_{H^1(\Omega)} \leq Ch^{-1}$ , which indicates that the error may increase as  $h \rightarrow 0$ .  $\blacksquare$

We note that for the case  $k = 1$ , Theorem 5.5 yields  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h + \eta)$ . In fact a similar result for  $k = 1$  can be obtained using less restrictions on the numerical integration. We state the result in the following corollary.

**Corollary 5.7** *Let  $u \in C^2(\bar{\Omega})$ ,  $a_{ij} \in C^1(\bar{\Omega})$ , and  $c \in C(\bar{\Omega})$ . Suppose the subspace  $V_h$ , with  $k = 1$ , satisfies assumptions **A1-A5** and the quadrature schemes*



satisfy **QA 4.1**, **QA 4.4**, and (4.12) only for  $i \in N_h''$ . Then, for  $\eta$  small enough, there is a positive constant  $C$ , independent of  $u$ ,  $\eta$ ,  $\tau$ , and  $h$ , such that

$$\|u - u_h^*\|_{H^1(\Omega)} \leq C(h + \eta + \tau)\|u\|_{W^{2,\infty}(\Omega)}.$$

The proof of this result can be obtained by slightly modifying the proofs of Lemma 5.4 and Theorem 5.5; we do not present the complete proof here.

## 6 Construction of numerical integration formula

In this section, we will derive numerical integration rules that satisfy the assumption **QA 4.2**, i.e., the condition (4.12) for  $k = 1$  and  $k = 2$  in two dimensions. We note that we have illustrated the conditions (4.12) in Remark 4.7.

To approximate the integral  $\int_{\omega_i}^s \varrho(x) dx$ , we seek a  $p$ -point quadrature rule  $Q_c^i(\varrho)$  on  $\omega_i$ , of the form

$$Q_c^i(\varrho) \equiv \sum_{l=1}^p \zeta_{c,l}^i \varrho(y_{c,l}^i), \quad \varrho \in C^0(\bar{\omega}_i) \text{ and } y_{c,l}^i \in \bar{\omega}_i, \quad (6.1)$$

that satisfies (4.12) with  $f_{\omega_i}^s$  replaced by  $Q_c^i$ .

**The case  $k = 1$ :** Recall that in this case, the shape functions  $\{\phi_i\}_{i \in N_h}$  reproduce polynomials of degree  $k = 1$ . We will find the weights  $\zeta_{c,l}^i$  and the integration points  $y_{c,l}^i$  in (6.1) such that (4.12), i.e., (4.19) and (4.20), are satisfied with  $f_{\omega_i}^s$  replaced by  $Q_c^i(\cdot)$ . Suppose we have at our disposal a quadrature rule

$$Q_B^i(g) \equiv \int_{\partial\omega_i \cap \Gamma} g(s) ds, \quad g \in C^0(\bar{\omega}_i) \quad (6.2)$$

that accurately approximates the boundary integral  $\int_{\partial\omega_i \cap \Gamma} g(s) ds$ . We start with an accurate  $p$ -point quadrature rule  $Q^i(\varrho)$  on  $\omega_i$  of the form

$$Q^i(\varrho) \equiv \sum_{l=1}^p \varrho(y_l^i) \zeta_l^i. \quad (6.3)$$

We then define for  $1 \leq l \leq p$ ,

$$\begin{aligned} y_{c,l}^i &= y_l^i \\ \zeta_{c,l}^i &= \zeta_l^i + \theta_1^i \zeta_l^i \frac{\partial \phi_i}{\partial x_1}(y_l^i) + \theta_2^i \zeta_l^i \frac{\partial \phi_i}{\partial x_2}(y_l^i), \end{aligned} \quad (6.4)$$

and choose  $\theta_1^i$  and  $\theta_2^i$  such that (4.19) and (4.20) are satisfied, i.e.,

$$\begin{aligned} Q_c^i\left(\frac{\partial \phi_i}{\partial x_1}\right) &= \int_{\partial\omega_i \cap \Gamma} n_1 \phi_i ds = Q_B^i(n_1 \phi_i) \\ Q_c^i\left(\frac{\partial \phi_i}{\partial x_2}\right) &= \int_{\partial\omega_i \cap \Gamma} n_2 \phi_i ds = Q_B^i(n_2 \phi_i). \end{aligned}$$

This yields the linear system

$$\begin{bmatrix} Q^i\left(\left(\frac{\partial\phi_i}{\partial x_1}\right)^2\right) & Q^i\left(\frac{\partial\phi_i}{\partial x_1}\frac{\partial\phi_i}{\partial x_2}\right) \\ Q^i\left(\frac{\partial\phi_i}{\partial x_2}\frac{\partial\phi_i}{\partial x_1}\right) & Q^i\left(\left(\frac{\partial\phi_i}{\partial x_2}\right)^2\right) \end{bmatrix} \begin{bmatrix} \theta_1^i \\ \theta_2^i \end{bmatrix} = \begin{bmatrix} Q_B^i(n_1\phi_i) - Q^i\left(\frac{\partial\phi_i}{\partial x_1}\right) \\ Q_B^i(n_2\phi_i) - Q^i\left(\frac{\partial\phi_i}{\partial x_2}\right) \end{bmatrix} \quad (6.5)$$

The components  $\theta_1^i$  and  $\theta_2^i$  of the solution of the above system are used in the definition of  $\zeta_{c,l}^i$  (see (6.4)), and consequently, the resulting  $Q_c^i(\varrho)$  satisfies the condition (4.12). We note that for  $i \in N_h''$ , the RHS of (6.5) does not contain the terms  $Q_B^i(n_1\phi_i)$  and  $Q_B^i(n_2\phi_i)$ . We further note that  $Q_c^i(\varrho)$  could be viewed as a corrected form of  $Q^i(\varrho)$ , such that  $Q_c^i(\varrho)$  satisfies the condition (4.12); we will often refer to  $Q_c^i(\varrho)$  as the *corrected numerical integration formula for  $k = 1$* . We note that  $Q_c^i(\varrho)$  for  $k = 1$  in the one dimensional case was derived in [6].

**Remark 6.1** To discuss the solvability of the system (6.5), we define a weighted inner product in  $\mathbb{R}^p$  by

$$\langle u, v \rangle_w \equiv \sum_{l=1}^p u_l v_l \zeta_l^i, \quad \forall u = (u_1, \dots, u_p) \text{ and } v = (v_1, \dots, v_p) \in \mathbb{R}^p$$

Let  $V_1 = \left(\frac{\partial\phi_i}{\partial x_1}(y_1^i), \dots, \frac{\partial\phi_i}{\partial x_1}(y_p^i)\right)$  and  $V_2 = \left(\frac{\partial\phi_i}{\partial x_2}(y_1^i), \dots, \frac{\partial\phi_i}{\partial x_2}(y_p^i)\right)$ , then the coefficient matrix of the linear system (6.5) is

$$\begin{bmatrix} \langle V_1, V_1 \rangle_w & \langle V_1, V_2 \rangle_w \\ \langle V_2, V_1 \rangle_w & \langle V_2, V_2 \rangle_w \end{bmatrix}, \quad (6.6)$$

which the Gramm matrix of the vectors  $V_1$  and  $V_2$  with respect to the inner product  $\langle \cdot, \cdot \rangle_w$ . This Gramm matrix is positive when  $V_1$  and  $V_2$  are linearly independent. Suppose  $p \geq 2$  and let there be two integration points  $y_m^i$  and  $y_n^i$  in the set of integration points  $\{y_l^i\}_{l=1}^p$  such that the vectors  $\nabla\phi_i(y_m^i)$  and  $\nabla\phi_i(y_n^i)$  are linearly independent, then it is easy to show that the vectors  $V_1$  and  $V_2$  are linearly independent. ■

**The case  $k = 2$ :** We recall that in this case, the shape functions  $\{\phi_i\}_{i \in N_h}$  reproduce polynomials of degree  $k = 2$ . We will find  $\zeta_{c,l}^i$  and  $y_{c,l}^i$  in (6.1) such that (4.12), with  $f_{\omega_i}^s$  replaced by  $Q_c^i$ , is satisfied for  $k = 2$ .

Suppose in addition to the quadrature rule  $Q_B^i(g)$  (see (6.2)), we also have at our disposal a quadrature rule

$$Q_F^i(f) \equiv \int_{\omega_i}^l f(x) dx$$

that accurately approximates the integral  $\int_{\omega_i}^l f(x) dx$ . As in the case  $k = 1$ , we start with an accurate  $p$ -point quadrature rule  $Q^i(\varrho)$  (see (6.3)). We note that we could choose  $Q^i(\cdot)$  to be same as  $Q_F^i(\cdot)$ . Suppose  $\mathcal{B} = \{\tilde{p}_m\}_{m=1}^6$  be a basis

for  $\tilde{\mathcal{P}}^1$  (recall that  $\dim \tilde{\mathcal{P}}^1 = 6$ ; a basis of  $\tilde{\mathcal{P}}^1$  is given in Remark 4.7). Then for  $k = 2$ , the condition (4.12), with  $f_{\omega_i}^s$  replaced by  $Q_c^i(\cdot)$ , is equivalent to

$$Q_c^i(\tilde{p}_m \cdot \nabla \phi_i) = Q_B^i(\tilde{p}_m \cdot \vec{n} \phi_i) - Q_F^i(\nabla \cdot \tilde{p}_m \phi_i), \quad \text{for } 1 \leq m \leq 6. \quad (6.7)$$

We now define, for  $1 \leq l \leq p$ ,

$$\begin{aligned} y_{c,l}^i &= y_l^i \\ \zeta_{c,l}^i &= \zeta_l^i + \sum_{n=1}^6 \theta_n^i \zeta_l^i(\tilde{p}_n \cdot \nabla \phi_i)(y_l^i), \end{aligned} \quad (6.8)$$

where  $\{\theta_n^i\}_{n=1}^6$  are chosen such that (6.7) is satisfied. We first note that from the definition of  $Q_c^i(\cdot)$  in (6.1), with  $y_{c,l}^i, \zeta_{c,l}^i$  as defined above, we get for  $1 \leq m \leq 6$ ,

$$\begin{aligned} Q_c^i(\tilde{p}_m \cdot \nabla \phi_i) &= Q^i(\tilde{p}_m \cdot \nabla \phi_i) + \sum_{l=1}^p \sum_{n=1}^6 \theta_n^i \zeta_l^i(\tilde{p}_n \cdot \nabla \phi_i)(y_l^i) (\tilde{p}_m \cdot \nabla \phi_i)(y_l^i) \\ &= Q^i(\tilde{p}_m \cdot \nabla \phi_i) + \sum_{n=1}^6 \theta_n^i Q^i\left((\tilde{p}_n \cdot \nabla \phi_i) (\tilde{p}_m \cdot \nabla \phi_i)\right) \end{aligned}$$

Therefore (6.7) is equivalent to the linear system

$$\begin{aligned} \sum_{n=1}^6 \theta_n^i Q^i\left((\tilde{p}_n \cdot \nabla \phi_i) (\tilde{p}_m \cdot \nabla \phi_i)\right) &= Q_B^i(\tilde{p}_m \cdot \vec{n} \phi_i) \\ &\quad - Q_F^i(\nabla \cdot \tilde{p}_m \phi_i) - Q^i(\tilde{p}_m \cdot \nabla \phi_i), \quad \text{for } 1 \leq m \leq 6. \end{aligned} \quad (6.9)$$

We use the solution  $\{\theta_n^i\}_{n=1}^6$  of the above linear system in the definition of  $\zeta_{c,l}^i$  (see (6.8)), and consequently,  $Q_c^i(\varrho)$  will satisfy the condition (4.12). We will often refer to  $Q_c^i(\varrho)$  as the *corrected numerical integration formula for  $k = 2$* . We note that solving the linear system (6.9) could be facilitated by considering the basis  $\mathcal{B} = \{\tilde{p}_m\}_{m=1}^6 = \{(1, 0), (x_1 - x_{i1}, 0), (x_2 - x_{i2}, 0), (0, 1), (0, x_1 - x_{i1}), (0, x_2 - x_{i2})\}$ , where  $x_i = (x_{i1}, x_{i2})$  is the particle associated with  $\omega_i$ .

**Remark 6.2** To discuss the solvability of the system (6.9), we define a weighted inner product for  $\mathbb{R}^p$  by

$$\langle u, v \rangle_w \equiv \sum_{l=1}^p u_l v_l \zeta_l^i, \quad \forall u = (u_1, \dots, u_p) \text{ and } v = (v_1, \dots, v_p) \in \mathbb{R}^p$$

Let  $V_n = \left( [\tilde{p}_n \cdot \nabla \phi_i](y_1^i), \dots, [\tilde{p}_n \cdot \nabla \phi_i](y_p^i) \right)$ ,  $1 \leq n \leq 6$ , then the coefficient matrix of the linear system (6.9) is

$$\begin{bmatrix} \langle V_1, V_1 \rangle_w & \langle V_1, V_2 \rangle_w & \cdots & \langle V_1, V_6 \rangle_w \\ \langle V_2, V_1 \rangle_w & \langle V_2, V_2 \rangle_w & \cdots & \langle V_2, V_6 \rangle_w \\ \vdots & \vdots & \ddots & \vdots \\ \langle V_6, V_1 \rangle_w & \langle V_6, V_2 \rangle_w & \cdots & \langle V_6, V_6 \rangle_w \end{bmatrix}$$

which is exactly the Gramm matrix of the vectors  $V_n$ ,  $1 \leq n \leq 6$  with respect to the inner product  $\langle \cdot, \cdot \rangle_w$ . This Gramm matrix is positive if the vectors  $\{V_l\}_{l=1}^6$  are linearly independent. We note that  $p \geq 6$  is a necessary condition for the linear independence of the vectors  $\{V_l\}_{l=1}^6$ . We further mention that the positivity of the Gramm matrix is subtle. When  $\omega_i$  is a square, our computations show that the Gramm matrix is positive when  $Q^i$  is the  $4 \times 4$  Gauss rule on  $\omega_i$ , but it has a zero eigenvalue when  $Q^i$  is the  $3 \times 3$  Gauss rule. ■

We now give a brief sketch of the derivation of the numerical integration rule  $Q_c^i(\cdot)$ , in 1 dimension (i.e., when  $d = 1$ ) with  $\omega_i = (\alpha_i, \beta_i)$ , such that (4.12) for  $k = 2$  is satisfied with  $f_{\omega_i}^s$  replaced by  $Q_c^i$ . We will use the one dimensional quadrature rule in our numerical examples in the next section.

As before, we start with the quadrature rules  $Q_F^i(\cdot)$  and  $Q^i(\cdot)$ ; we recall that both the rules could also be same. We first note that, for  $d = 1$ , the “boundary integral” term in (4.11) is  $v\phi_i|_{\alpha_i}^{\beta_i}$ ; thus we do not need the quadrature rule  $Q_B^i(\cdot)$ . We further note that  $\tilde{\mathcal{P}}^1 = \mathcal{P}^1$  and therefore  $m = \dim \mathcal{P}^1 = 2$ . Thus (6.7), for  $d = 1$ , is written as

$$\begin{cases} Q_c^i(\phi_i'(x)) = \phi_i(x)|_{\alpha_i}^{\beta_i} \\ Q_c^i([(x - x_i)\phi_i'(x)]) = (x - x_i)\phi_i(x)|_{\alpha_i}^{\beta_i} - Q_F^i(\phi_i) \end{cases} \quad (6.10)$$

We now define  $y_{c,l}^i, \zeta_{c,l}^i$  (compare with (6.8)) as

$$\begin{cases} y_{c,l}^i = y_l^i \\ \zeta_{c,l}^i = \zeta_l^i + \theta_1^i \zeta_l^i \phi_i'(y_l^i) + \theta_2^i \zeta_l^i [(y_l^i - x_i)\phi_i'(y_l^i)], \end{cases} \quad (6.11)$$

where  $\theta_1^i, \theta_2^i$  are chosen such that (6.10) is satisfied. Using  $Q_c^i(\cdot)$ , with  $y_{c,l}^i, \zeta_{c,l}^i$  as defined above, in (6.10) yields the linear system for  $\theta_1^i, \theta_2^i$ , namely,

$$\begin{aligned} & \begin{bmatrix} Q^i(\phi_i'^2(x)) & Q^i[(x - x_i)\phi_i'^2(x)] \\ Q^i[(x - x_i)\phi_i'^2(x)] & Q^i[(x - x_i)^2\phi_i'^2(x)] \end{bmatrix} \begin{bmatrix} \theta_1^i \\ \theta_2^i \end{bmatrix} \\ &= \begin{bmatrix} \phi_i(x)|_{\alpha_i}^{\beta_i} - Q^i(\phi_i') \\ (x - x_i)\phi_i(x)|_{\alpha_i}^{\beta_i} - Q^i[(x - x_i)\phi_i'(x)] - Q_F^i(\phi_i) \end{bmatrix}. \end{aligned} \quad (6.12)$$

## 7 Numerical Results

We present numerical examples to illuminate the results obtained in Section 5. Let  $\Omega = (0, 1)$  and we consider the Neumann problem with non-constant coefficients, namely,

$$\begin{aligned} -(au')' + cu &= f, \quad x \in \Omega \\ a(0)u'(0) &= 1, \quad a(1)u'(1) = 2e, \end{aligned}$$

where  $a(x) = 1 + x^3$ ,  $c(x) = 1 + \sin^2 x$ , and  $f(x) = e^x(\sin^2 x - x^3 - 3x^2)$ . The exact solution of the problem is  $u(x) = e^x$ .

To approximate the solution  $u(x)$  of the above problem by the meshless method (4.6), we first define the shape functions of the finite dimensional space  $V_h$ . For a given non-negative integer  $k$  and a positive real number  $R$ , let  $\phi(x)$  be the basic RKP shape function with compact support  $[-R, R]$  satisfying

$$\sum_{i \in \mathbb{Z}} i^l \phi(x - i) = x^l, \quad \forall x \in \mathbb{R} \text{ and } l = 0, 1, \dots, k. \quad (7.1)$$

We mention that there exists  $\phi(x)$  satisfying (7.1) when  $R \geq (k + 1)/2$  (see e.g., [3]). Consider a positive integer  $N$  and for  $h = 1/N$ , we consider the index set

$$N_h = \{-[R], \dots, 0, 1, \dots, N, \dots, N + [R]\},$$

where  $[R]$  is the integer part of  $R$ . For each  $i \in N_h$ , we define the RKP shape functions

$$\phi_i(x) \equiv \phi\left(\frac{x}{h} - i\right), \quad x \in \Omega.$$

Then  $\text{supp } \phi_i = [\alpha_i, \beta_i] = [ih - Rh, ih + Rh] \cap [0, 1]$ . Defining the set of particles  $X_h = \{x_i = ih, i \in N_h\}$ , it can be easily shown that  $\{\phi_i\}_{i=1}^{N_h}$  reproduce polynomials of degree  $k$ , i.e.,

$$\sum_{i \in N_h} x_i^l \phi_i(x) = x^l, \quad \forall x \in \Omega \text{ and } l = 0, 1, \dots, k.$$

Moreover, recalling the definitions of the index sets  $N'_h$  and  $N''_h$ , we have

$$N'_h = \{-[R], \dots, [R], N - [R], \dots, N + [R]\} \text{ and } N''_h = \{[R] + 1, \dots, N - [R] - 1\}$$

We note that the function  $\phi(x)$  has been constructed following the ideas mentioned in Remark 3.3 (using  $h = 1$ ,  $x_j = j \in \mathbb{Z}$ , and  $i = 0$ , i.e.,  $\phi(x) = \phi_0^1(x)$ ), where we have used the cubic spline weight function for  $w(x)$  with compact support  $[-R, R]$ ; for the definition of cubic spline weight function, we refer to [3, 4]. We further note that the cubic spline weight function is symmetric in  $[-R, R]$ , and consequently the associated shape functions  $\phi_i(x)$ ,  $i \in N''_h$ , are also symmetric in  $[\alpha_i, \beta_i]$ .

**The case  $k = 1$**

The basic shape function  $\phi(x)$  was constructed with  $R = 1.8$ . For  $i \in N_h$ , we consider the standard  $p$ -point Gaussian integration rule on  $[\alpha_i, \beta_i]$ , namely,

$$Q_g^i(f) \equiv \sum_{l=1}^p f(y_l^i) \zeta_l^i, \quad \forall f \in C(\omega_i),$$

where  $\{y_l^i : 1 \leq l \leq p\}$  are the Gaussian integration points in  $[\alpha_i, \beta_i]$  and  $\{\zeta_l^i : 1 \leq l \leq p\}$  are the associated weights. It is well known that the points  $y_l^i$  are symmetrically placed in the interval  $[\alpha_i, \beta_i]$ ; also the weights  $\zeta_l^i$  are symmetric, i.e.,  $\zeta_s^i = \zeta_{p+1-s}^i$ ,  $s = 1, 2, \dots, p$ .

Recall that  $\phi_i(x)$ , for  $i \in N_h''$ , is symmetric, and consequently,  $\phi_i'(x)$ ,  $i \in N_h''$ , is anti-symmetric in  $[\alpha_i, \beta_i]$  about the mid-point. Therefore, we get


$$Q_g^i(\phi_i') = 0, \quad \forall i \in N_h''. \quad (7.2)$$

Thus the numerical integration rule  $Q_g^i$  satisfies the condition (4.12), i.e., the discrete Green's formula, for  $i \in N_h''$  (see also (4.21)). We used  $Q_g^i$ , with  $p = 8, 16, 32$ , and  $64$  to compute  $\gamma_{ij}^*$ ,  $\sigma_{ij}^*$ , and  $l_i^*$  in the variational problem (4.6). We note that in the one dimensional case, evaluation of the boundary integrals is trivial. We further note that  $\eta$  decreases as  $p$  increases. We have computed the solution  $u_h^*$  of (4.6) and have presented the error  $\|u - u_h^*\|_{H^1(\Omega)}$  for various values of  $h$  in Table 1. We also presented the log-log graph of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$  in Figure 1. It is clear that for  $p = 16, 32$  and  $64$ , the error  $\|u - u_h^*\|_{H^1(\Omega)}$  first decreases and then ‘‘levels off’’, which suggests that  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h + \eta)$ . This illuminates the result of the Corollary 5.7.

**Table 1.** MM with  $k = 1$ . Standard Gaussian integration rule

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	1.9883E-02	3.4312E-03	3.4040E-03	3.3908E-03
1/20	2.3933E-02	1.7751E-03	1.7851E-03	1.7427E-03
1/40	2.6763E-02	9.1993E-04	9.8673E-04	8.8411E-04
1/80	2.8425E-02	4.9672E-04	6.4523E-04	4.4625E-04
1/160	2.9324E-02	3.0333E-04	5.3111E-04	2.2612E-04
1/320	2.9791E-02	2.2761E-04	5.0200E-04	1.1769E-04
1/640	3.0029E-02	2.0272E-04	4.9631E-04	6.7007E-05
1/1280	3.0150E-02	1.9519E-04	4.9581E-04	4.6261E-05

**Table 1.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 1$ . Standard  $p$ -point Gaussian integration rule, with  $p = 8, 16, 32$  and  $64$ , was used in the MM. These rules satisfy the discrete Green's formula in the interior for  $k = 1$ , i.e., the assumption (7.2).



**Figure 1.** The loglog plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduce polynomials of degree  $k = 1$ . Standard  $p$ -point Gaussian rule is used, which satisfy the assumption (7.2).

We now show that the condition (4.12) on the underlying quadrature rule is a necessary condition for the result presented in Theorem 5.5, i.e.,  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h + \eta)$ . We consider a non-symmetric Gaussian integration rule that does not satisfy the condition (4.12), i.e., does not satisfy the discrete Green's formula. For  $i \in N_h$ , we consider the mapping  $T_i : [\alpha_i, \beta_i] \rightarrow [\alpha_i, \beta_i]$  given by

$$z = T_i(y) = y + \frac{0.2}{\beta_i - \alpha_i} \left[ \left( y - \frac{\alpha_i + \beta_i}{2} \right)^2 - \left( \frac{\beta_i - \alpha_i}{2} \right)^2 \right]$$

Therefore, for a smooth function  $f$ , we have

$$\int_{\alpha_i}^{\beta_i} f(z) dz = \int_{\alpha_i}^{\beta_i} f(T_i(y)) T_i'(y) dy.$$

The integral on the RHS of the above equality could be approximated by the Gaussian rule  $Q_g^i$  to obtain an integration rule on  $[\alpha_i, \beta_i]$  to approximate the integral  $\int_{\alpha_i}^{\beta_i} f(z) dz$ , namely,


$$Q_{ng}^i(f) \equiv \sum_{l=1}^p f(y_{nl}^i) \zeta_{nl}^i, \quad (7.3)$$

where  $y_{nl}^i = T_i(y_l^i)$  and  $\zeta_{nl}^i = T_i'(y_l^i) \zeta_l^i$ . We will refer to  $Q_{ng}^i$  as a  $p$ -point non-symmetric Gaussian integration rule on  $[\alpha_i, \beta_i]$ . It is well known that the algebraic precision of  $Q_g^i$  is  $2p - 1$ ; we can show that the algebraic precision of  $Q_{ng}^i$  is  $p - 1$ .

**Table 2.** MM with  $k = 1$ . Non-symmetric Gaussian integration rule.

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	4.5373E-01	8.5286E-03	3.4098E-03	3.3908E-03
1/20	1.1694E+00	2.0281E-02	2.0558E-03	1.7425E-03
1/40	2.7436E+00	4.4534E-02	2.8166E-03	8.8444E-04
1/80	6.3125E+00	9.2759E-02	5.9275E-03	4.5245E-04
1/160	1.4474E+01	1.8844E-01	1.2420E-02	2.7798E-04
1/320	3.2278E+01	3.7713E-01	2.5442E-02	3.5063E-04
1/640	6.9243E+01	7.4415E-01	5.1451E-02	6.6803E-04
1/1280	1.4414E+02	1.4379E+00	1.0329E-01	1.3328E-03

**Table 2.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 1$ . Non-symmetric  $p$ -point Gaussian integration rule, with  $p = 8, 16, 32$  and  $64$ , was used in the MM. These integration rules *do not* satisfy the discrete Green's formula for  $k = 1$ .



**Figure 2.** The loglog plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 1$ . Non-symmetric  $p$ -point Gaussian integration rules were used, which *do not* satisfy the discrete Green's formula for  $k = 1$ .

We use the non-symmetric Gaussian integration rule  $Q_{ng}^i$ , with  $p = 8, 16, 32$ , and  $64$ , to compute  $\gamma_{ij}^*$ ,  $\sigma_{ij}^*$ , and  $l_i^*$  in the variational problem (4.6). We computed the solution  $u_h^*$  of (4.6) and presented the error  $\|u - u_h^*\|_{H^1(\Omega)}$  for various values of  $h$  in Table 2. We also presented the log-log plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with



respect to  $h$  in Figure 2. It is clear that  $\|u - u_h^*\|_{H^1(\Omega)}$  increases as  $h$  decreases; for  $p = 32$  and  $64$ , the error first decreases and then increases. In all the cases, the error  $\|u - u_h^*\|_{H^1(\Omega)}$  behaves like  $\mathcal{O}(h^{-1})$ , as indicated in Corollary 5.6 and Remark 5.3.

Now following the ideas presented in Section 6, we will correct the non-symmetric Gaussian integration rule  $Q_{ng}^i(\cdot)$  (given in (7.3)), such that the corrected numerical integration rule (see (6.1))

$$Q_c^i(\varrho) \equiv Q_{ng,c}^i(\varrho) = \sum_{l=1}^p \varrho(y_{c,l}^i) \zeta_{c,l}^i$$

satisfies the condition (4.12). We note that for  $d = 1$ , the condition (4.12) for  $k = 1$  is

$$Q_{ng,c}^i(\phi'_i) = \phi(\beta_i) - \phi(\alpha_i). \quad (7.4)$$

For  $1 \leq i \leq p$ , we consider

$$y_{c,l}^i = y_{nl}^i \quad \text{and} \quad \zeta_{c,l}^i = \zeta_{nl}^i + \theta^i \zeta_{nl}^i \phi'_i(y_{nl}^i),$$

with

$$\theta^i = \frac{\phi_i(\beta_i) - \phi_i(\alpha_i) - \sum_{l=1}^p \phi'_i(y_{nl}^i) \zeta_{nl}^i}{\sum_{l=1}^p \phi_i'^2(y_{nl}^i) \zeta_{nl}^i}.$$


Then it can be shown, following the ideas in Section 6 for  $d = 1$ , that  $Q_{ng,c}^i(\cdot)$  satisfies the condition (4.12), i.e., (7.4). However, we note that unlike the standard Gaussian integration rule  $Q_g^i(\cdot)$ , the integration points for the quadrature rule  $Q_{ng,c}^i(\cdot)$  are not symmetrically placed in  $[\alpha_i, \beta_i]$ . The expression for the corrected numerical integration rule for  $d = 1$  was also derived in [6] for a slightly different situation. We will refer to  $Q_{ng,c}^i(\cdot)$  as the *corrected non-symmetric Gaussian integration rule for  $k = 1$* .

We now use the corrected integration rule  $Q_{ng,c}^i(\cdot)$  to compute  $\gamma_{ij}^*$  in problem (4.6); the terms  $\sigma_{ij}^*$  and  $l_i^*$  in (4.6) are computed using the integration rule  $Q_{ng}^i$  (uncorrected). We computed the solution  $u_h^*$  of (4.6) and have presented the error  $\|u - u_h^*\|_{H^1(\Omega)}$ , for various values of  $h$  in Table 3. We also presented the log-log plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$  in Figure 3. It is clear that  $\|u - u_h^*\|_{H^1(\Omega)}$  levels off as  $h$  decreases; the error first decreases and then levels off for  $p = 16, 32$ , and  $64$ . This suggests that  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h + \eta)$ .

**Table 3.** MM with  $k = 1$ . Corrected non-symmetric Gaussian integration rule for  $k = 1$ .

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	5.6887E-03	3.4243E-03	3.4004E-03	3.3907E-03
1/20	4.8897E-03	1.7538E-03	1.7735E-03	1.7424E-03
1/40	4.9036E-03	9.8324E-04	9.5897E-04	8.8351E-04
1/80	5.0486E-03	7.3643E-04	5.9529E-04	4.4492E-04
1/160	5.1580E-03	7.0595E-04	4.6444E-04	2.2331E-04
1/320	5.2221E-03	7.2101E-04	4.2812E-04	1.1201E-04
1/640	5.2564E-03	7.3597E-04	4.2001E-04	5.6271E-05
1/1280	5.2742E-03	7.4520E-04	4.1871E-04	2.8400E-05

**Table 3.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 1$ . Corrected non-symmetric Gaussian integration rule for  $k = 1$ , with  $p = 8, 16, 32$  and  $64$ , was used in the MM. The integration rules satisfy the discrete Green's formula for  $k = 1$ .



**Figure 3.** The loglog plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduce polynomials of degree  $k = 1$ . Corrected  $p$ -point non-symmetric Gaussian rules for  $k = 1$  were used, which satisfy the discrete Green's formula for  $k = 1$ .

### The case $k = 2$


The basic shape function  $\phi(x)$ , satisfying (7.1) with  $k = 2$ , was constructed with  $R = 2.2$ . Let  $Q^i(\cdot) = Q_{ng}^i(\cdot)$  be the  $p$ -point non-symmetric Gaussian integration rule on  $[\alpha_i, \beta_i]$ , as given in (7.3). We consider the associated corrected

non-symmetric Gaussian integration rule  $Q_c^i(\cdot)$  for  $k = 2$ ;  $Q_c^i(\cdot)$  satisfies the discrete Green's formula (4.12) for  $k = 2$ ,  $d = 1$ , i.e., it satisfies (6.10). The integration points  $\{y_{c,l}^i\}_{l=1}^p$  and the associated weights  $\{\zeta_{c,l}^i\}_{l=1}^p$  of  $Q_c^i(\cdot)$  are given by (6.11) with  $y_l^i = y_{n,l}^i$  and  $\zeta_l^i = \zeta_{n,l}^i$  for  $1 \leq l \leq p$ . We mention that  $\theta_1^i, \theta_2^i$  in (6.11) are obtained from the solution of (6.12), with  $Q_F^i(\phi_i) = Q_{ng}^i(\phi_i)$ . We used the corrected non-symmetric Gaussian integration rule  $Q_c^i(\cdot)$  (for  $k = 2$ ) to compute the terms  $\gamma_{ij}^*$  in the variational problem (4.6). The terms  $\sigma_{ij}^*$  and  $l_i^*$  were computed using the non-symmetric Gaussian integration rule (uncorrected)  $Q_{ng}^i(\cdot)$ . We computed the solution  $u_h^*$  of (4.6) and have presented the values of  $\|u - u_h^*\|_{H^1(\Omega)}$ , for various values of  $h$  in Table 4. We also presented the log-log plot of the ratio  $\frac{\|u - u_h^*\|_{H^1(\Omega)}}{h}$  with respect to  $h$  in Figure 4. It is clear that  $u_h^*$  converges to  $u$ , the solution of (2.3), as  $h$  becomes smaller. The Figure 4 also indicates that  $\|u - u_h^*\|_{H^1(\Omega)} = O[h(h + \eta)]$ , illuminating the result of Theorem 5.5 for  $k = 2$ .

**Table 4.** MM with  $k = 2$ . Corrected non-symmetric Gaussian rule for  $k = 2$ .

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	6.8778E-03	1.3037E-03	7.0255E-04	6.8283E-04
1/20	4.4966E-03	8.3073E-04	2.1238E-04	1.7378E-04
1/40	2.5677E-03	4.8134E-04	8.4965E-05	4.4034E-05
1/80	1.3736E-03	2.5972E-04	4.1096E-05	1.1258E-05
1/160	7.1071E-04	1.3494E-04	2.0824E-05	3.0523E-06
1/320	3.6153E-04	6.8783E-05	1.0558E-05	9.6564E-07
1/640	1.8234E-04	3.4724E-05	5.3252E-06	3.8283E-07
1/1280	9.1564E-05	1.7446E-05	2.6751E-06	1.7719E-07

**Table 4.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 2$ . Corrected  $p$ -point non-symmetric Gaussian integration rule for  $k = 2$ , with  $p = 8, 16, 32$  and  $64$ , was used in the MM. The integration rules satisfy the discrete Green's formula for  $k = 2$ .




**Figure 4.** The loglog plot of the ratio  $\frac{\|u - u_h^*\|_{H^1(\Omega)}}{h}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduce polynomials of degree  $k = 2$ . Corrected  $p$ -point non-symmetric Gaussian rules for  $k = 2$  were used, which satisfy the discrete Green's formula for  $k = 2$ .

We will now show the effect of quadrature on  $\|u - u_h^*\|_{H^1(\Omega)}$ , when the quadrature does not satisfy the condition (4.12) for  $k = 2$ . We first computed  $\gamma_{ij}^*$  in (4.6) using the corrected  $p$ -point non-symmetric integration rule for  $k = 1$  (see  $Q_{ng,c}^i(\cdot)$  given before). We note that this quadrature rule satisfies only the first condition in (6.10). The terms  $\sigma_{ij}^*$  and  $l_i^*$  were computed using the  $p$ -point non-symmetric gaussian integration rule  $Q_{ng}^i(\cdot)$ . The error  $\|u - u_h^*\|_{H^1(\Omega)}$  for various values of  $h$  and the associated log-log plot is given in Table 5 and Figure 5 respectively. These results indicate that  $\|u - u_h^*\|_{H^1(\Omega)} = \mathcal{O}(h + \eta)$  and  $u_h^* \rightharpoonup u$  as  $h \rightarrow 0$ . Finally, we used  $Q_{ng}^i(\cdot)$  to compute  $\gamma_{ij}^*$  in (4.6);  $Q_{ng}^i(\cdot)$  does not satisfy any of the conditions in (6.10). The terms  $\sigma_{ij}^*$  and  $l_i^*$  were again computed using  $Q_{ng}^i(\cdot)$ . The error  $\|u - u_h^*\|_{H^1(\Omega)}$  for various values of  $h$  and the associated log-log plot is given in Table 6 and Figure 6 respectively. It is clear that the error  $\|u - u_h^*\|_{H^1(\Omega)}$  diverges as  $h$  decreases; in fact  $\|u - u_h^*\|_{H^1(\Omega)}$  behaves like  $\mathcal{O}(h^{-1})$  as suggested by Corollary 5.6 for  $k = 2$ .

**Table 5.** MM with  $k = 2$ . Corrected non-symmetric Gaussian rule for  $k = 1$ .

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	3.8161E-02	6.3683E-03	1.8719E-03	6.9042E-04
1/20	4.0997E-02	1.2019E-02	1.9499E-03	2.2130E-04
1/40	3.9586E-02	1.5615E-02	2.0005E-03	1.5928E-04
1/80	3.8040E-02	1.7666E-02	2.0156E-03	1.6104E-04
1/160	3.7035E-02	1.8762E-02	2.0196E-03	1.6434E-04
1/320	3.6472E-02	1.9330E-02	2.0205E-03	1.6610E-04
1/640	3.6174E-02	1.9619E-02	2.0208E-03	1.6699E-04
1/1280	3.6022E-02	1.9764E-02	2.0208E-03	1.6743E-04

**Table 5.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 2$ . Corrected  $p$ -point non-symmetric Gaussian integration rule for  $k = 1$  (not corrected for  $k = 2$ ), with  $p = 8, 16, 32$  and  $64$ , was used in the MM. The integration rules do not satisfy the discrete Green's formula for  $k = 2$ .




**Figure 5.** The loglog plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduce polynomials of degree  $k = 2$ . Corrected  $p$ -point non-symmetric Gaussian rules for  $k = 1$  were used, which do not satisfy the discrete Green's formula for  $k = 2$ .

**Table 6.** Non-symmetric(no correction) Gaussian rule:  $k=2$ .

$h$	$\ u - u_h^*\ _{H^1(\Omega)}$			
	8 points	16 points	32 points	64 points
1/10	9.4346E-01	4.6767E-02	7.5982E-03	7.6165E-04
1/20	2.0155E+00	4.5294E-02	2.1562E-02	1.0294E-03
1/40	4.3714E+00	9.2717E-02	5.0461E-02	2.3620E-03
1/80	9.7615E+00	2.8436E-01	1.0883E-01	5.0628E-03
1/160	2.1885E+01	7.1121E-01	2.2511E-01	1.0474E-02
1/320	4.7659E+01	1.6259E+00	4.5406E-01	2.1309E-02
1/640	1.0044E+02	3.6444E+00	8.9677E-01	4.3013E-02
1/1280	2.0680E+02	8.2709E+00	1.7231E+00	8.6554E-02

**Table 6.** The  $H^1$  norm of the error,  $\|u - u_h^*\|_{H^1(\Omega)}$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduced polynomials of degree  $k = 2$ . Non-symmetric  $p$ -point Gaussian integration rule, with  $p = 8, 16, 32$  and  $64$ , was used in the MM. The integration rules do not satisfy the discrete Green's formula for  $k = 2$ .



**Figure 6.** The loglog plot of  $\|u - u_h^*\|_{H^1(\Omega)}$  with respect to  $h$ , where  $u = e^x$  and  $u_h^*$  is the solution of the MM, employing shape functions that reproduce polynomials of degree  $k = 2$ . Non-symmetric  $p$ -point Gaussian integration rules were used, which do not satisfy the discrete Green's formula for  $k = 2$ .

## References

- [1] S. N. Atluri and S. Shen. *The Meshless Local Petrov Galerkin Method*. Tech. Sci. Press, 2002.

- [2] S. N. Atluri and T. Zhu. A new meshless local petrov-galerkin (mlpg) approach in computational mechanics. *Computational Mechanics*, 22:117–127, 1998.
- [3] I. Babuška, U. Banerjee, and J. Osborn. Survey of meshless and generalized finite element methods: a unified approach. *Acta Numerica*, 12:1–125, 2003.
- [4] I. Babuška, U. Banerjee, and J. Osborn. On the approximability and the selection of particle shape functions. *Numer. Math.*, 96:601–640, 2004.
- [5] I. Babuška, U. Banerjee, J. Osborn, and Q. Li. Quadrature for meshless methods. *Int. J. Numer. Meth. Engng.*, 76:1434–1470, 2008. to be published.
- [6] I. Babuška, U. Banerjee, J. Osborn, and Q. Zhang. Effect of numerical integration on meshless methods. *Comput. Methods Appl. Mech. Engrg.*, 198:2886–2897, 2009.
- [7] S. Beissel and T. Belytschko. Nodal integration of the element-free Galerkin method. *Comput. Methods Appl. Mech. Engrg.*, 139:49–74, 1996.
- [8] T. Belytschko, Y. Krongauz, D. Organ, M. Fleming, and P. Krysl. Meshless methods: An overview and recent developments. *Comp. Meth. Appl. Mech. Engrg.*, 139:3–47, 1996.
- [9] T. Belytschko, Y. Y. Lu, and L. Gu. Element-free Galerkin methods. *Int. J. Numer. Meth. Engng.*, 37:229–256, 1994.
- [10] S. C. Brenner and L. R. Scott. *The Mathematical Theory of Finite Element Methods*. Springer-Verlag, New York, 2007.
- [11] A. Carpinteri, G. Ferro, and G. Ventura. The partition of unity quadrature in meshless methods. *Int. J. Numer. Meth. Engng.*, 54:987–1006, 2002.
- [12] A. Carpinteri, G. Ferro, and G. Ventura. The partition of unity quadrature in element free crack modelling. *Computers and Structures*, 81:1783–1794, 2003.
- [13] J.-S. Chen, C.-T. Wu, S. Yoon, and Y. You. A stabilized conformal nodal integration for a Galerkin mesh-free method. *Int. J. Numer. Meth. Engng.*, 50:435–466, 2001.
- [14] J.-S. Chen, S. Yoon, and C.-T. Wu. Non-linear version of stabilized conforming nodal integration Galerkin mesh-free methods. *Int. J. Numer. Meth. Engng.*, 53:2587–6515, 2002.
- [15] P. G. Ciarlet. *The Finite Element Method for Elliptic Problems*. North-Holland, Amsterdam, 1978.
- [16] S. De and K. J. Bathe. The method of finite spheres. *Computational Mechanics*, 25:329–345, 2000.

- [17] S. De and K.J. Bathe. The method of finite squares with improved numerical integration. *Computers and Structures*, 79:2183–2196, 2001.
- [18] J. Dolbow and T. Belytschko. Numerical integration of the Galerkin weak form in meshfree methods. *Computational Mechanics*, 23:219–230, 1999.
- [19] T. -P. Fries and H. -G. Matthies. Classification and Overview of Meshfree Methods. Technical report, Technical University Braunschweig, Brunswick, Germany, 2004.
- [20] M. Griebel and M. A. Schweitzer. A particle-partition of unity method. ii. efficient cover construction and reliable integration. *SIAM J. Sci. Comput.*, 23:1655–1682, 2002.
- [21] W. Han and X. Meng. Error analysis of the reproducing kernel particle method. *Comput. Methods Appl. Mech. Engrg*, 190:6157–6181, 2001.
- [22] S. Li and W. K. Liu. Meshfree and particle methods and their application. *Applied Mechanics Review*, 55:1–34, 2002.
- [23] S. Li and W. K. Liu. *Meshfree Particle Methods*. Springer, 2004.
- [24] W. K. Liu, S. Jun, and Y. F. Zhang. Reproducing kernel particle methods. *Int. J. Numer. Meth. Fluids*, 20:1081–1106, 1995.
- [25] J. M. Melenk. On approximation in meshless methods. In J. Blowey and A. Craig, editors, *Frontiers in Numerical Analysis, Durham 2004*. Springer Verlag, 2005.
- [26] E. M. Stein. *Singular Integrals and Differentiability Properties of Functions*. Princeton Univ. Press, 1970.
- [27] G. Strang. Variational crimes in the finite element method. In A. K. Aziz, editor, *Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations*, pages 698–710. Academic Press, 1972.